

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 960 937 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

01.12.1999 Bulletin 1999/48

(21) Application number: 97939229.7

(22) Date of filing: 09.09.1997

(51) Int. Cl.⁶: **C12N 15/22**, A61K 38/17,
A61K 31/70, A01K 67/027,
C07K 14/475, G01N 33/577,
A61K 39/395

(86) International application number:
PCT/JP97/03167

(87) International publication number:
WO 98/11216 (19.03.1998 Gazette 1998/11)

(84) Designated Contracting States:

**AT BE CH DE DK ES FI FR GB GR IE IT LI NL PT
SE**

(30) Priority: 11.09.1996 JP 26356596
08.08.1997 JP 22722097

(71) Applicant:

**SUMITOMO PHARMACEUTICALS COMPANY,
LIMITED
Osaka-shi, Osaka-fu 541 (JP)**

(72) Inventors:

- **KIMURA, Toru**
Kusatsu-shi, Shiga 525 (JP)
- **KIKUCHI, Kaoru**
Takarazuka-shi, Hyogo 665 (JP)

(74) Representative:

**VOSSIUS & PARTNER
Siebertstrasse 4
81675 München (DE)**

(54) **NOVEL SEMAPHORIN GENE: SEMAPHORIN Y**

(57) The present invention provides Semaphorin Y inhibiting neurite outgrowth, and a gene therefor, as well as other Semaphorins hybridizing to said Semaphorin Y gene, modified proteins or partial peptides of said Semaphorin Y, antibodies against said Semaphorin Y, anti-sense nucleotides against said Semaphorin Y gene, and the use of such substances as pharmaceutical or diagnostic agents or laboratory reagents. The present invention further provides a method of screening for Semaphorin Y antagonists employing said Semaphorin Y, Semaphorin Y antagonists obtained by said screening method, pharmaceutical agents comprising such antagonists, and transgenic animals involving said Semaphorin Y.

EP 0 960 937 A1

Description

TECHNICAL FIELD

5 [0001] The present invention relates to Semaphorin Y, a novel Semaphorin belonging to the Semaphorin family, and use of Semaphorin Y for pharmaceutical or diagnostic agents or laboratory reagents. More particularly, it relates to Semaphorin Y inhibiting neurite outgrowth, and a gene therefor, as well as other Semaphorins hybridizing to said Semaphorin Y gene, modified proteins or partial peptides of said Semaphorin Y, antibodies against said Semaphorin Y, anti-sense nucleotides against said Semaphorin Y gene, antagonists of said Semaphorin Y, transgenic animals, and their
10 use as pharmaceutical or diagnostic agents or laboratory reagents.

BACKGROUND ART

15 [0002] It is widely known that a central nervous system (CNS)-neuron in higher organisms such as human is not capable of regeneration once injured. Therefore, one who has received an injury on his (her) spinal cord due to, for example, a traffic accident, is compelled to spend the rest of his (her) life in a hemiplegic state. On the contrary, it is known that a peripheral nervous system (PNS)-neuron retains a vigorous regeneration ability even in those higher organisms, and therefore, neurons in a limb, when disconnected, can gradually regenerate with a concomitant recovery of their function.

20 [0003] In the early nineteen-eighties, a group of Aguayo *et al.* found that when PNS-neuron is experimentally grafted into an injured CNS-neuron in a higher organism, axon growth of CNS-neuron is induced. This observation demonstrates that CNS-neuron in higher organisms which had been generally considered not to have a regeneration ability can regenerate if a suitable environment is provided (*Nature*, **284**, 264-265 (1980), *Science*, **214**, 931-933 (1981)). That report suggests a possibility that in CNS of higher organisms, there may exist a factor, namable "CNS-neuron regeneration inhibitor", which inhibits the regeneration of CNS-neuron, and that a release from such inhibition may
25 allow the regeneration of CNS-neurons. This suggestion paved the way for a CNS-neuron regeneration therapy.

30 [0004] In 1988, a group of Schwab *et al.* demonstrated that there existed such CNS-neuron regeneration inhibitor among proteins derived from CNS myelin. They also succeeded in purifying, though partially, a protein having said CNS-neuron regeneration inhibition activity, and named this protein fraction NI35/250 (*Annu. Rev. Neurosci.*, **16**, 565-595 (1993)), although no one has succeeded in its isolation, identification and gene cloning yet. In addition, they immunized animals with the partial purified NI35/250, and succeeded in obtaining an antibody (IN-1) having a neutralizing activity. This antibody is capable of recognizing a band for NI35/250 in Western blotting, and capable of staining, in an immunostaining, the region to which NI35/250 is supposed to be distributed. Furthermore, they demonstrated that administration of this antibody to an animal experimentally received an injury on its spinal cord has promoted regeneration of axons in spinal cord, though partially, within 2-3 weeks, and restored its function within 2-3 months (*Nature*,
35 **343**, 269-272 (1990), *Nature*, **378**, 498-501 (1995)). These findings are of great value, because they experimentally demonstrated that there existed a CNS-neuron regeneration inhibitor as suggested by Aguayo *et al.* (*supra*) and that CNS-neuron can be regenerated by inhibiting the activity of said inhibitor. The above antibody is, however, directed not to human but to rat NI35/250, and exhibits a low stability and specificity. In addition, although regeneration of CNS-neuron was observed as described above by administering said antibody, its effect was so partial and incomplete that not
40 all of the motor functions could be restored. It is, therefore, believed essential in solving these problems to identify the gene for NI35/250 or corresponding CNS-neuron regeneration inhibitor, and, based on knowledges of molecular biology, neuroscience and the like, develop an antagonist more effectively inhibiting the CNS-neuron regeneration inhibition activity, or develop a method for inhibiting the expression of the gene for said regeneration inhibitor.

45 [0005] Apart from the above, the nervous system, whether it is central or peripheral, requires formation of a complicated neural network among neurons or between neurons and peripheral receivers or effectors during development, that is, in the stage of embryo or fetus, in order to precisely carry out its principal functions, *i.e.*, to transfer and process the information. To establish the neural network, an ingenious mechanism is necessary, which precisely guides a growing neurite to the target site locating remote therefrom.

50 [0006] It has been hitherto believed that a factor which positively controls the neurite outgrowth, such as neurite growth promoter and neurite growth attractant may play a major role in the formation of the neural network. However, it is now being demonstrated by recent studies on the mechanism of the network formation that the opposite factor, that is, a negative factor having an outgrowth inhibition activity is important for an accurate guidance (*Cell*, **78**, 353-356 (1994)).

55 [0007] A representative factor having such an outgrowth inhibition activity is a protein called "Semaphorin". Semaphorin firstly discovered is Fasciclin IV found in grasshopper. Collapsin (latterly named Collapsin I) was subsequently discovered in chick (*Cell*, **75**, 217-227 (1993); *Neuron*, **9**, 831-845 (1992)). To date, more than 10 genes belonging to the Semaphorin family have been reported in a wide range of species covering insects such as drosophila and beetle, human, and viruses (*Cell*, **81**, 471-474 (1995)). These Semaphorins characteristically contain in their amino acid

sequences similar structures called semaphorin domains each consisting of about 500 amino acids (*Neuron*, 14, 941-948 (1995); *Cell*, 75, 1389-1399 (1993)). However, the homologies of the primary amino acid sequences in semaphorin domains among these Semaphorin genes are 80-20%, and not necessarily high.

- 5 [0008] Of these Semaphorins, functions have been verified for only a few, including, for example, Fasciclin IV of grasshopper, Semaphorins I and II of drosophila, Collapsin of chick, and Semaphorin III which corresponds to Collapsin in mammals. All of these Semaphorins are known to inhibit neurite outgrowth or synapsis formation. In particular, Semaphorin III has been reported to have an activity collapsing in a short time the growth cone of cultured neuron (growth-cone collapse activity) *in vitro* (*Neuron*, 14, 941-948 (1995); *Neuron*, 14, 949-959 (1995); *Cell*, 81, 631-639 (1995); *Cell*, 75, 1389-1399 (1993); *Cell*, 75, 217-227 (1993); *Neuron*, 9, 831-845 (1992)).
- 10 [0009] Although it is now being demonstrated, as described above, that known Semaphorins have a growth-cone collapse activity and a neurite outgrowth inhibition activity during development, and play a role in giving an accurate guidance to neuron, it is not evident at present whether or not their Semaphorins exert some function not only during development but also in the adult, and less evident whether or not Semaphorins play a role as a CNS-neuron regeneration inhibitor. Of course, since known Semaphorins have been shown to be a negative guidance factor inhibiting neurite outgrowth, it would not be unreasonable to consider said Semaphorins as a candidate for a CNS-neuron regeneration inhibitor (*Nature*, 378, 439-440 (1995)). However, it has been shown by *in vitro* experiments that Semaphorin III (Sema III), only one Semaphorin of higher organisms of which function has been analyzed, exerts its neurite-outgrowth inhibition activity on a sensory neuron and sympathetic neuron both of which are peripheral, but not on a retinal neuron which is central (*Cell*, 75, 217-227 (1993)). In addition, Northern analysis on the distribution of Sema III expression in the adult conducted by the present inventors has revealed that it is expressed mainly in peripheral tissues (see Reference example 2 below). It is therefore hardly believed that Sema III having such features has a function as a CNS-neuron regeneration inhibitor.

PROBLEM TO BE SOLVED BY THE INVENTION

- 25 [0010] The present invention aims to provide Semaphorin Y, a novel Semaphorin belonging to the Semaphorin family, and a gene therefor, and to provide pharmaceutical agents for neural diseases, in particular for regeneration of CNS-neuron, and related diagnostic agents or laboratory reagents. More specifically, the present invention aims to provide Semaphorin Y inhibiting neurite outgrowth and a gene therefor, as well as other Semaphorins hybridizing to said Semaphorin Y gene, modified proteins or partial peptides of said Semaphorin Y, antibodies against said Semaphorin Y, antisense nucleotides against said Semaphorin Y gene, and use of such substances as pharmaceutical or diagnostic agents or laboratory reagents. The present invention further aims to provide a method of screening for Semaphorin Y antagonists employing said Semaphorin Y, Semaphorin Y antagonists obtained by said screening method, pharmaceutical agents comprising such antagonists, and transgenic animals involving Semaphorin Y.

MEANS OF SOLVING THE PROBLEM

- 35 [0011] In order to provide pharmaceutical agents for neural diseases, in particular for regeneration of CNS-neuron, and related diagnostic agents or laboratory reagents, the present inventors have planned to identify a novel Semaphorin which has not yet been cloned. In particular, the present inventors have paid their attention to the similarity between the *in vitro* activities of the above-described NI35/250 and Semaphorin, i.e., to the fact that NI35/250 has a growth-cone collapse activity and a neurite-growth inhibition activity *in vitro* (*J. Neurosci.*, 8, 2381-2393 (1988); *Science*, 259, 80 (1993)), while known Semaphorins similarly possess a neurite-growth inhibition activity, and particularly Semaphorin III has also a growth-cone collapse activity. This suggested to the inventors the possibility that unknown Semaphorins which have not yet been identified may include the one inhibiting regeneration of CNS-neuron. Specifically, the present inventors' idea was that Semaphorin, which is characterized in that 1) it is widely expressed throughout the CNS of adult where regeneration of neuron (or neurite outgrowth) is inhibited, but 2) it is poorly expressed in other tissues such as peripheral tissues in the adult, has not been identified yet, and if one can identify a new unknown Semaphorin having such characteristics, the Semaphorin might be involved in inhibition of regeneration of CNS-neuron.
- 40 [0012] First of all, the inventors have closely searched DNA database on the basis of the amino acids sequence relatively well conserved among previously reported Semaphorin genes. Specifically, a DNA sequence has been searched through EST (Expressed Sequence Tags) database, which is a gene not expressed in peripheral tissues but expressed in the postnatal brain and which encodes an amino acid sequence relatively well conserved among Semaphorins. As a consequence, a DNA fragment R59527 was identified, which encodes, as a partial sequence, a sequence consisting of seven amino acids: Gln (or Arg)-Asp-Pro-Tyr-Cys-Ala (or Gly)-Trp. The R59527 gave a sequence information as to only 238 bases, and furthermore only several percent thereof could be translated into an amino acid sequence common to those of known Semaphorins. In addition, the reading frame could not be determined because of the presence of sequence not definitely determined in R59527. It was, therefore, utterly impossible at that stage to con-

clude that the base sequence of R59527 is part of a novel Semaphorin. We have, however, finally succeeded in cloning a novel Semaphorin gene by carrying out the following procedures: synthesizing DNA primers on the basis of that sequence information; conducting PCR with said primers using cDNAs prepared from a human hippocampal cDNA library as templates to obtain a novel DNA fragment (SEQ ID NO: 7) consisting of 170 bases; labeling the DNA fragment with ³²P to synthesize a DNA probe; and screening rat and human cDNA libraries with that probe. We named this novel Semaphorin "Semaphorin Y".

[0013] Subsequent analysis revealed that Semaphorin Y is a novel Semaphorin at which we aimed, since it was widely expressed in CNS in the adult, whereas among peripheral tissues the expression could be observed only in limited tissues.

[0014] Semaphorin Y of the present invention having such characteristics appears to be involved in inhibition of regeneration of CNS-neuron in the adult. Semaphorin Y may be used to screen for Semaphorin Y antagonists, and antagonists identified in such screening system are expected to promote regeneration of CNS-neuron. Similarly, antisense DNAs or RNAs against Semaphorin Y gene are also expected to promote regeneration of CNS-neuron as the above antagonists do.

[0015] In addition, in view of the fact that Semaphorin Y of the present invention inhibits neurite outgrowth, it may be used as a therapeutic or diagnostic agent for pains or immune diseases such as atopic dermatitis, by administering it to peripheral tissues, which results in the inhibition of neurite outgrowth of PNS-neuron. Furthermore, Semaphorin Y is a novel Semaphorin belonging to the Semaphorin family of which expression distribution is unconventionally characteristic as described above, and also has a characteristic in that it does not contain any Ig domains commonly found among hitherto reported Semaphorins of vertebrates. Semaphorin Y may, therefore, serve as an important research material or a laboratory reagent.

[0016] The present invention has been completed on the basis of the above findings.

[0017] Thus, the gist of the present invention is as follows:

(1) a gene encoding the following protein (a) or (b):

- (a) Semaphorin Y protein comprising the amino acid sequence shown in SEQ ID NO: 3 or 6,
- (b) a protein which comprises an amino acid sequence wherein one or more amino acids are deleted, substituted and/or added in the amino acid sequence shown in SEQ ID NO: 3 or 6, and which protein inhibits neurite outgrowth;

(2) a gene comprising the following DNA (a) or (b):

- (a) Semaphorin Y DNA comprising the base sequence shown in SEQ ID NO: 1, 2, 4, or 5,
- (b) DNA which hybridizes under stringent conditions to DNA comprising the base sequence shown in SEQ ID NO: 1, 2, 4, or 5, and which encodes a protein inhibiting neurite outgrowth;

(3) a gene comprising DNA which hybridizes under stringent conditions to DNA comprising the base sequence shown in SEQ ID NO: 7, and which encodes a protein having a semaphorin domain;

(4) a protein obtained by expressing the gene of any one of the above items (1) to (3);

(5) a gene comprising DNA which encodes a protein comprising an amino acid sequence in which one or more amino acids are deleted, substituted and/or added in the protein shown in SEQ ID NO: 3 or 6, wherein said protein promotes neurite outgrowth;

(6) a protein obtained by expressing the gene of the above item (5);

(7) DNA which is cloned from a human cDNA library or a human genomic library, and which hybridizes under stringent conditions to DNA comprising at least part of DNA consisting of the base sequence shown in SEQ ID NO: 1 or 4;

(8) an expression plasmid which expresses either the gene of any one of the above items (1) to (3) and (5), or DNA of the above item (7);

(9) a transformant transformed with the expression plasmid of the above item (8);

(10) a process for producing a recombinant protein, which process comprises culturing the transformant of the above item (9), and recovering the recombinant protein expressed;

(11) a peptide comprising at least six amino acids of the protein of the above item (4) or (6);

(12) a peptide of the above item (11) which promotes neurite outgrowth;

(13) a peptide of the above item (11) characterized in that it contains aspartic acid residue at position 198 of the amino acid sequence shown in SEQ ID NO: 6 or an amino acid residue corresponding to the position of said aspartic acid residue;

(14) an antisense nucleotide, or chemically modified variant thereof, which is directed against a segment compris-

ing at least eight or more bases in the gene of any one of the above items (1) to (3), or in DNA of the above item (7);
 (15) an antisense nucleotide or chemically modified variant thereof of the above item (14), characterized in that it
 inhibits expression of the protein of the above item (4);
 (16) an antibody against the protein of the above item (4) or (6), or against the peptide of any one of the above items
 5 (11) to (13);
 (17) a pharmaceutical agent comprising, as an active ingredient, the gene of any one of the above items (1) to (3)
 and (5), DNA of the above item (7), the protein of the above item (4) or (6), the peptide of any one of the above
 items (11) to (13), the antisense nucleotide or chemically modified variant thereof of the above item (14) or (15), or
 the antibody of the above item (16);
 10 (18) a method of screening for Semaphorin Y antagonists, characterized in that it employs the protein of the above
 item (4);
 (19) Semaphorin Y antagonist obtained by the screening method of the above item (18);
 (20) Semaphorin Y antagonist of the above item (19) which comprises the protein of the above item (6), the peptide
 of any one of the above items (11) to (13), or the antibody of the above item (16);
 15 (21) a CNS-neuron regeneration promoter, characterized in that it contains at least one of the antisense nucle-
 otides or chemically modified variants thereof of the above item (14) or (15), or Semaphorin Y antagonists of the
 above item (19) or (20);
 (22) a neurite outgrowth inhibitor for PNS-neuron, characterized in that it contains at least one of the proteins of the
 above item (4); and
 20 (23) a transgenic animal in which either the gene of any one of the above items (1) to (3) and (5), or DNA of the
 above item (7) has been artificially inserted into its chromosome, or has been knocked out.

MODE FOR CARRYING OUT THE INVENTION

25 **[0018]** The 1st embodiment of the present invention is a gene which encodes Semaphorin Y comprising the amino
 acid sequence shown in SEQ ID NO: 3 or 6, or a gene encoding a protein which comprises an amino acid sequence
 wherein one or more amino acids are deleted, substituted and/or added in the amino acid sequence of the above Sem-
 aphorin Y, and which protein inhibits neurite outgrowth. The 2nd embodiment of the present invention is Semaphorin Y
 30 gene comprising the base sequence shown in SEQ ID NO: 1, 2, 4, or 5, or a gene which hybridizes under stringent con-
 ditions to such Semaphorin Y gene and which encodes a protein inhibiting neurite outgrowth. These genes are
 explained below in order.

1) Gene Encoding Semaphorin Y (Semaphorin Y Gene)

35 **[0019]** Of the above-mentioned genes, "a gene which encodes Semaphorin Y protein comprising the amino acid
 sequence shown in SEQ ID NO: 3" or "Semaphorin Y gene comprising the base sequence shown in SEQ ID NO: 1 or
 2" is a gene encoding the rat Semaphorin Y of the present invention, while "a gene which encodes Semaphorin Y pro-
 tein comprising the amino acid sequence shown in SEQ ID NO: 6" or "Semaphorin Y gene comprising the base
 40 sequence shown in SEQ ID NO: 4 or 5" is a gene encoding the human Semaphorin Y of the present invention. Among
 these genes, those shown in SEQ ID NOs: 2 and 5 correspond open reading frames for rat and human types of Sem-
 aphorin Y, respectively. Such genes may be cloned, as described in Example 1, by screening a cDNA library derived
 from CNS tissues using a probe (for example, a DNA probe having the base sequence shown in SEQ ID NO: 7) pre-
 pared on the basis of the sequence of "R59527" found in EST database. Particular techniques for such cloning may be
 found in the standard texts such as "Molecular Cloning, 2nd ed.", Cold Spring Harbor Laboratory Press (1989). The
 45 base sequence of the cloned DNA may also be determined by conventional methods, for example, using a sequencing
 kit commercially available.

[0020] Alternatively, after publication of the base sequence of rat and human Semaphorin Y cDNAs of the present
 invention, one skilled in the art can also easily clone the full-length genes encoding rat and human types of Semaphorin
 Y by using part of said cDNA as a probe, without using cloning methods as described above.

50 2) Gene Encoding Modified Protein of Semaphorin Y

[0021] Of the above-mentioned genes, "a gene encoding a protein which comprises an amino acid sequence wherein
 one or more amino acids are deleted, substituted and/or added in the amino acid sequence of the above Semaphorin
 55 Y, and which protein inhibits neurite outgrowth" refers to a gene encoding a so-called "modified proteins" of Semaphorin
 Y which inhibits neurite outgrowth. Those skilled in the art may easily obtain a gene encoding such protein, for example,
 by site-directed mutagenesis (*Methods in Enzymology*, 100, 448- (1983)) or PCR method (*Molecular Cloning*, 2nd ed.,
 Chapter 15, Cold Harbor Laboratory Press (1989); "PCR A Practical Approach", IRL Press, 200-210 (1991)). In this

context, the number of amino acid residues to be deleted, substituted and/or added is to be such a number that permits the deletion, substitution and/or addition by well-known methods such as site-directed mutagenesis described above.

[0022] For the purpose of the present invention, the phrase "inhibiting neurite outgrowth" means that the protein has the collapse activity on growth cone of neuron, or that the protein has the neurite-outgrowth inhibition activity. These activities may be measured with a test substance such as an expression product of DNA encoding Semaphorin Y or modified protein thereof, for example, in the following manner:

[0023] Since Semaphorin Y is a membrane protein, it exists in the cell membrane of the cells transformed with Semaphorin Y gene. The activities of the above test substance may, therefore, easily be measured by using, as a test material, the membrane fraction of the transformed cells.

[0024] Examples of activity measurement include measurement of collapse activity on growth cone of neuron (M. Igarashi *et al.*, *Science*, vol. 259, pp. 77-79 (1993)), or measurement of neurite-outgrowth inhibition activity (e.g., J. A. Davies *et al.*, *Neuron*, vol. 2, pp. 11-20 (1990) and M. Bastmeyer, *J. Neurosci.*, vol. 11, pp. 626-640 (1991)). A method of measuring the growth-cone collapse activity is described in detail in literature (M. Igarashi *et al.*, *Science*, vol. 259, pp. 77-79 (1993)). Briefly, the measurement may be carried out by a method in which cells expressing a test substance such as Semaphorin Y is homogenized, and the homogenate containing the cell membrane fraction or the purified membrane fraction is used (E. C. Cox *et al.*, *Neuron*, vol. 2, pp. 31-37 (1990)), or by a method in which a protein extracted from the membrane fraction is reconstituted in a liposome and the liposome is used as a test material (C. E. Bandtlow, *Science*, vol. 259, pp. 80-84 (1993)). In order to measure the growth-cone collapse activity in practice using these materials, a test substance such as Semaphorin Y, for example, in one of the forms as describe above is added to neurons cultured under conventional conditions (e.g., "Culturing, Nerve Cells" edited by Banker *et al.*, MIT Press (1991)) in a container coated with a substance promoting the neurite outgrowth and the growth-cone formation such as laminin, collagen, polylysine or polyornithine. After the addition, when a sufficient time has passed to occur collapse of growth cone (typically from 30 minutes to one hour after the addition), those neurons are fixed with 1% glutaraldehyde or the like, and the number of the growth cones which have been collapsed is counted under a microscope. In this measurement, it is important that another sample is used as a control, which is prepared from cells not expressing the test substance such as Semaphorin Y according to the completely same procedures as those used for the test substance-expressing cells. Typically, normalization of the samples is conducted on the basis of the total amounts of protein included within the samples. To measure the neurite-outgrowth inhibition activity, part of the surface of a micropore filter or a culture container made of glass or plastics is coated with a test substance such as Semaphorin Y prepared as described above, and the activity is indicated, for example, by the inability of neurons cultured under conventional conditions to adhere to the coated area, or by a remarkable decrease in the rate of neurite outgrowth on the coated area, or by the inability of invasion of growing neurites from the outside of the coated area into the coated area because of its stopping on the border between the coated and non-coated areas or its avoidance from the coated area. When a cluster of cells expressing a test substance is co-cultured with neurons in a collagen gel, the inability of outgrowing neurite to enter the cluster of cells expressing the test substance may also be used as an indicator (A. Sophia *et al.*, *Cell*, vol. 81, 621-629 (1995)).

[0025] Both neurons of CNS and PNS may be used as the cells for the above activity measurements. As described in the section "BACKGROUND ART", CNS in adult mammals naturally contains a large amount of regeneration (outgrowth) inhibitor. It is, therefore, extremely difficult to measure *in vivo* an inhibitory effect on neurite outgrowth of CNS-neuron, and such inhibitory effect is usually measured by an *in vitro* method as described above. Since these *in vitro* methods each have individual characteristics, it is preferred to use more than one method to confirm the activity. Although preferred neurons used for a measurement of the activity are CNS-neurons such as motor neurons in spinal cord or motor cortex, PNS-neurons in superior cervical ganglion and dorsal root ganglion may also be used because NI35/250 known as a CNS-neuron regeneration inhibitor has proved to have effects such as neurite-growth inhibition and growth-cone collapse activities also on such PNS-neurons (*J. Cell Biol.*, 106, 1281-1288 (1988); *Science*, 259, 80-83 (1993)).

[0026] Specific examples of the modified proteins of this embodiment are described below.

[0027] Based on the structural comparison of known Semaphorins, most of the conserved amino acids are located in the semaphorin domain, suggesting that these conserved amino acids are essential for expression of the activity of Semaphorins. Furthermore, the present inventors have found that a modified Sema III protein in which aspartic acid residue at position 198 in its semaphorin domain has been substituted with glycine did not have the growth-cone collapse activity (see Reference example 1 below). Accordingly, the aspartic acid at position 198 of Sema III is believed essential for expression of the activity. The amino acid residues corresponding to this position are highly conserved in known Semaphorins, and they are all aspartic acid with a few exceptions in which glutamic acid is located at this position. It is, therefore, believed that the amino acid residue at this position is also essential for expression of the activity of Semaphorins other than Sema III. In Semaphorin Y of the present invention, the amino acid residue corresponding to the position 198 of Sema III is presumed to be aspartic acid at position 197 in the amino acid sequence shown in SEQ ID NO: 3, or aspartic acid at position 198 in the amino acid sequence of human Semaphorin Y shown in SEQ ID NO: 6.

[0028] Considering the above information, it is desirable to make the above-described deletions, substitutions and/or additions of amino acids at positions other than those conserved among Semaphorins, so as to retain the activity of Semaphorin Y in modified proteins. Particularly, it is desirable not to modify the aspartic acid at position 197 in rat Semaphorin Y shown in SEQ ID NO: 3 and the aspartic acid at position 198 in human Semaphorin Y. In order to substitute amino acids conserved among Semaphorins while retaining the activity of Semaphorin Y, it is desirable to substitute an amino acid having a similar side chain for the amino acid residue to be substituted. By substituting such amino acid having a similar side chain for a conserved amino acid, it may be possible to produce a modified protein which has an enhanced activity of Semaphorin Y. Such modified protein having the enhanced activity is highly suitable as a neurite-outgrowth inhibitor for PNS-neuron as will be described hereinafter in the section of the 22nd embodiment of the present invention.

[0029] In the above-noted embodiment, "a conserved amino acid" refers to an amino acid located at a position at which more than 50% of Semaphorin genes shown in Fig. 2 of *Cell*, 75, 1389-1399 (1993) or Fig. 1 of *Neuron*, 14, 941-948 (1995) share the same amino acid.

3) DNA Hybridizing Under Stringent Conditions To Semaphorin Y Gene

[0030] Of the above-mentioned DNAs, "a gene which hybridizes under stringent conditions to Semaphorin Y gene and which encodes a protein inhibiting neurite outgrowth" refers to a gene such as Semaphorin Y gene derived from a mammal, which hybridizes under stringent conditions to rat or human Semaphorin Y gene comprising the base sequence shown in SEQ ID NO: 1, 2, 4, or 5.

[0031] As used herein, "a gene which hybridizes under stringent conditions" refers to such a gene that hybridizes to rat or human Semaphorin Y gene, for example, when subjected to hybridization at a formamide concentration of about 45% (v/v) and a salt concentration of about 5x SSPE and at a temperature around 42°C, and washed at a salt concentration of about 2x SSPE and at a temperature around 42°C. Cloning of such genes may be achieved, for example, by screening cDNA or genomic libraries prepared from various animal tissues using all or part of DNA shown in SEQ ID NO: 1 or 4 as a probe. Such screening may be carried out by making reference to the standard texts such as "Molecular Cloning 2nd ed." (Cold Spring Harbor Laboratory Press (1989)).

[0032] Specific examples of the gene of this embodiment may include all the Semaphorin Y genes of mammal and avian. Between mammals or between mammal and avian, homologous genes have quite similar sequences, and usually more than 80%, in many cases more than 90%, of the base sequence are common to each other. All the mammal and avian Semaphorin Y genes, therefore, correspond to this embodiment. In other words, those genes which have a homology of 80% or above, and preferably of 90% or above, are included in this embodiment.

[0033] The 3rd embodiment of the present invention is a gene comprising DNA which hybridizes under stringent conditions to DNA comprising the base sequence shown in SEQ ID NO: 7, and which encodes a protein having a semaphorin domain.

[0034] In the above description, "DNA comprising the base sequence shown in SEQ ID NO: 7" refers to a fragment cloned by PCR using the sequence information of the DNA "R59527" which encodes, in part, a sequence consisting of seven amino acids well conserved among Semaphorins (Gln (or Arg)-Asp-Pro-Tyr-Cys-Ala (or Gly)-Trp), and the DNA fragment corresponds to a region from position 1574 to position 1743 in the base sequence of rat Semaphorin Y shown in SEQ ID NO: 1, or a region from position 1524 to position 1693 in the base sequence of human Semaphorin Y shown in SEQ ID NO: 4.

[0035] The "stringent conditions" refers to those conditions described above in the section of the 2nd embodiment of the present invention.

[0036] Cloning of these DNAs is achieved by, for example, hybridization with DNA of SEQ ID NO: 7, and specifically may be carried out, for example, according to the procedures described in *TINS*, 15, 319-323 (1992) and references cited therein, and more specifically according to the following procedures.

[0037] That is, the cloning may be achieved by screening cDNA or genomic libraries prepared from various animal tissues using DNA consisting of the base sequence shown in SEQ ID NO: 7 as a probe. The screening may be carried out according to, for example, the procedures as described in Example 1. Preferred cDNA libraries are those derived from an adult tissue of CNS, and cDNA libraries derived from hippocampus, corpus striatum, and cerebellum are more preferred. As described above, the conditions shown in Example 1 or those described in *TINS*, 15, 319-323 (1992) and references cited therein may be used for the hybridization.

[0038] The DNA of this embodiment is also "DNA which encodes a protein having a semaphorin domain". As used herein, "semaphorin domain" refers to a domain consisting of 300-600 amino acid residues more than 20% of which are identical to those amino acids constituting the semaphorin domain of any one of ten known Semaphorins (G-Sema I, T-Sema, I, D-Sema II, H-Sema III, C-Collapsin, Sem A, Sem B, Sem C, Sem D, Sem E) described in, for example, *Cell*, 75, 1389-1399 (1993) or *Neuron*, 14, 941-948 (1995). Those proteins having a semaphorin domain more than 30% of which amino acids are identical to those amino acids in any one of the known Semaphorins are particularly preferred.

The identity of amino acids is determined by comparison using, for example, DNASIS Ver. 2.0 (HITACH Software Engineering) under conditions of ktup=1 and cutoff=1. More preferred proteins are those in which ten or more cysteines, particularly twelve or more cysteines, of the thirteen cysteines conserved in semaphorin domains of the ten known Semaphorins (for example, those cysteines marked in Figure 1 on page 942 of *Neuron*, 14, 941-948 (1995)) are conserved.

[0039] Examples of such gene of this embodiment may include Semaphorin genes which hybridize under stringent conditions to DNA comprising the base sequence shown in SEQ ID NO: 7 and which contain semaphorin domains and exhibit the neurite-outgrowth inhibition activity, including all of the Semaphorin Y genes of mammal and avian.

[0040] The 4th embodiment of the present invention is a protein obtained by expressing the gene of any one of the above items (1) to (3).

[0041] Typical examples of protein included in this embodiment are rat Semaphorin Y comprising the amino acid sequence shown in SEQ ID NO: 3, and human Semaphorin Y comprising the amino acid sequence shown in SEQ ID NO: 6. The rat or human Semaphorin Y contains a signal sequence at its N-terminus and such signal sequence is presumed to correspond to a region from position 1 to position 23 of the amino acid sequence shown in SEQ ID NO: 3 or from position 1 to position 24 of the amino acid sequence shown in SEQ ID NO: 6, respectively. Since the signal sequence is removed by processing during its transfer to membrane, such mature forms of Semaphorin Y are also included in this embodiment.

[0042] Preparation of the proteins of this embodiment may be achieved, for example, by ligating a cloned rat Semaphorin Y cDNA into a known expression vector such as pET or pCDM8, and introducing it into appropriate host cells to express and produce Semaphorin Y. The host cells may be prokaryotic or eukaryotic. For example, *Escherichia coli* strains or animal cell lines are already conventionally used for such purpose and are commercially or publicly available. Examples of animal host cells include COS-1, COS-7, CHO cells and the like.

[0043] To transform appropriate animal host cells with an expression plasmid, a known procedure such as DEAE-dextran method (*Current Protocols in Molecular Biology*, F. M. Ausubel et al. ed., John Wiley & Sons (1987)) may be used. As confirmed in Example 6, Semaphorin Y exists in the cell membrane fraction which contains a sufficient amount of Semaphorin Y to be directly used in various assays. Therefore, various assays for activities of a protein of this embodiment may easily be conducted using a cell membrane fraction prepared from appropriate cells.

[0044] Furthermore, a protein of this embodiment may be purified by, for example, affinity purification using Semaphorin Y-recognizing antibodies described hereinafter in the section of the 16th embodiment of the present invention, or conventional column chromatography.

[0045] The 5th embodiment of the present invention is a gene encoding a protein which comprises an amino acid sequence wherein one or more amino acids are deleted, substituted and/or added in the rat or human Semaphorin Y shown in SEQ ID NO: 3 or 6 and which protein promotes neurite outgrowth. The 6th embodiment of the present invention is a protein obtained by expressing the gene of the 5th embodiment of the present invention.

[0046] In the genes of the above 5th embodiments, deletions, substitutions and/or additions may be introduced in the procedures similar to those used for a gene encoding a modified protein of the 1st embodiment of the present invention. Similarly, the promotion effect on neurite outgrowth may easily be measured, for example, by adding Semaphorin Y in an assay system for Semaphorin Y activity described above in the section of the 1st embodiment of the present invention and further adding thereto a test substance (i.e., a candidate modified Semaphorin Y protein). For details, see the descriptions in the section of the 18th embodiment of the present invention.

[0047] Specific examples of the proteins of the 6th embodiment may be modified proteins of which neurite-outgrowth inhibition activity has been eliminated. Such modified protein lacking the neurite-outgrowth inhibition activity is expected to exert the promotion effect on neurite-outgrowth, when it binds to receptors for Semaphorin Y or to Semaphorin Y itself, by inhibiting the binding of Semaphorin Y to the receptors. As described above in the section of the 1st embodiment of the present invention, it has been suggested that the active site of Semaphorin may be located in the semaphorin domain, and particularly, it may be located at aspartic acid at position 197 in rat Semaphorin Y or aspartic acid at position 198 in human Semaphorin Y. Accordingly, in order to eliminate the semaphorin Y activity from the modified protein, it is desirable to introduce the deletions, substitutions and/or additions to the conserved amino acids in said semaphorin domain, preferably to the aspartic acid at position 197 in rat Semaphorin Y or to the aspartic acid at position 198 in human Semaphorin Y. In such cases, those substitutions in which an amino acid having a side chain of a distinct nature is substituted for the original amino acid are desirable. Also in the cases of Semaphorin Y other than that from human or rat, modifications are preferably made on aspartic acid at this position, that is, on amino acid residue at the position which corresponds to position 197 in rat Semaphorin Y or to position 198 in human Semaphorin Y when the amino acid sequence of said Semaphorin Y is aligned with that of rat or human Semaphorin Y so as to give the maximum identity.

[0048] Since the proteins of the 6th embodiment of the present invention promote neurite outgrowth as described above, some of these proteins will serve as CNS-neuron regeneration promoters as described hereinafter in the section of the 21st embodiment.

[0049] The 7th embodiment of the present invention is DNA which is cloned from a human cDNA library or a human genomic library, and which hybridizes under stringent conditions to DNA comprising at least part of rat or human Semaphorin Y DNA shown in SEQ ID NO: 1 or 4, respectively.

[0050] Methods of cloning are described in detail in, for example, "Molecular Cloning 2nd ed.", Cold Spring Harbor Laboratory Press (1989), and specifically include, for example, methods employing hybridization or PCR. Although a preferred library used herein is a genomic library derived from human, a cDNA library derived from CNS-neuron in the adult may also be used. Those methods employing hybridization may be carried out according to, for example, *TINS*, 15, 319-323 (1992) and references cited therein. Those methods employing PCR may be carried out according to, for example, "PCR" edited by McPherson *et al.*, IRL Press (1991).

[0051] The DNAs thus cloned include not only the full length DNA but also its DNA fragments comprising more than 200 bases, or single-stranded forms (coding strands or complementary strands thereof) of said DNA fragments. Specific examples of DNA of the 7th embodiment of the present invention may include chromosomal DNAs containing 5' and/or 3' transcriptional control regions, noncoding sequences of exons, introns or the like, in addition to regions encoding amino acids. Such sequences which do not encode any amino acids are also quite useful, for example, in developing a medicine using antisense techniques described hereinafter.

[0052] The 8th embodiment of the present invention is an expression plasmid which expresses either the gene of the 1st, 2nd, 3rd or 5th embodiment, or DNA of the 7th embodiment of the present invention. The 9th embodiment of the present invention is a transformant transformed with the expression plasmid of the 8th embodiment. Furthermore, the 10th embodiment of the present invention is a process for producing a recombinant protein which process comprises culturing the transformant of the 9th embodiment and recovering the recombinant protein expressed. As described above in the section of the 4th embodiment of the present invention, methods of preparing an expression plasmid and a transformant, and methods of producing a recombinant protein, *per se*, are all well known to those skilled in the art.

[0053] The 11th embodiment of the present invention is a peptide comprising at least 6 amino acids of a protein of the 4th or 6th embodiment of the present invention. In this context, the limitation "at least 6 amino acids" is based on the fact that a minimal size of peptide capable of forming a stable structure consists of 6 amino acids, and preferred peptides are those consisting of 8 or more amino acids, more preferably of about 10-20 amino acids. A short peptide such as those consisting of about 10-20 amino acids can be synthesized on a peptide synthesizer, while a longer peptide can be obtained by preparing DNA through conventional genetic engineering, and expressing it in, for example, animal cells as described above. The peptide thus prepared can also be modified by conventional methods.

[0054] These peptides can be applied to pharmaceutical agents described hereinafter in the section of the 12th and 13th embodiments, and can also be used for producing antibodies.

[0055] The 12th embodiment of the present invention is a peptide of the 11th embodiment of the present invention which promotes neurite outgrowth. Such polypeptide may be prepared by the methods described above in the section of the 11th embodiment of the present invention. The promotion effect on neurite outgrowth can also easily be measured as described above in the section of the 5th embodiment of the present invention by adding Semaphorin Y to an activity assay system described above in the section of the 1st embodiment of the present invention and further adding thereto a test substance (*i.e.*, a candidate peptide of Semaphorin Y). For details, see the descriptions in the section of the 18th embodiment of the present invention.

[0056] Specific examples of these peptides may be peptides which have lost the neurite-outgrowth inhibition activity of Semaphorin Y. A peptide lacking Semaphorin Y activity is expected to exert its neurite-outgrowth promotion effect, when it binds to receptors for Semaphorin Y or to Semaphorin Y itself, by inhibiting the binding of Semaphorin Y to the receptors. Some of such peptides will serve as CNS-neuron regeneration promoters as described hereinafter in the section of the 21st embodiment.

[0057] The 13th embodiment of the present invention is a peptide of the 11th embodiment of the present invention, characterized in that it contains the aspartic acid residue at position 198 of the amino acid sequence shown in SEQ ID NO: 6 or an amino acid residue corresponding to the position of said aspartic acid residue. Such peptides may be prepared by the methods described above in the section of the 11th embodiment.

[0058] As described above in the section of the 1st embodiment of the present inventions, the aspartic acid residue at position 198 of human Semaphorin Y shown in SEQ ID NO: 6 (in the case of rat, the aspartic acid residue at position 197) seems essential for expression of the activity of Semaphorin Y. Since this amino acid residue may possibly be involved in the binding between Semaphorin Y and its receptors, a peptide of this embodiment containing this amino acid residue may interfere the neurite-outgrowth inhibition activity of Semaphorin Y by binding to receptors for Semaphorin Y or to Semaphorin Y itself, resulting in promotion of neurite outgrowth. Some of the peptides having such effect will serve as CNS-neuron regeneration promoters as described hereinafter in the section of the 21st embodiment. Such neurite-outgrowth promotion activity can easily be measured as described above in the section of the 5th embodiment of the present invention by adding Semaphorin Y to an activity assay system described in the section of the 1st embodiment of the present invention, and further adding thereto a test substance (*i.e.*, a candidate peptide of Semaphorin Y). For details, see the descriptions in the section of the 18th embodiment of the present invention.

[0059] In this embodiment, "an amino acid corresponding to the position of said aspartic acid" refers to an amino acid residue which is located at the position corresponding to position 198 in human Semaphorin Y, when the amino acid sequence of the protein of the 4th or 6th embodiment of the present invention is aligned with the amino acid sequence of human Semaphorin Y shown in SEQ ID NO: 6 so as to give the maximum identity. Accordingly, "a peptide characterized in that it contains an amino acid corresponding to the position of said aspartic acid" refers to a peptide which comprises such amino acid at the position corresponding to position 198 in human Semaphorin Y as well as flanking amino acids on either side thereof.

[0060] The 14th embodiment of the present invention is an antisense nucleotide, or chemically modified variant thereof, which is directed against a segment comprising at least eight or more bases in the gene of any one of the 1st to 3rd embodiments, or in DNA of the 7th embodiment of the present invention.

[0061] As used herein, "antisense nucleotide" refers to a so-called antisense oligonucleotide, antisense RNA, or antisense DNA, and it may be artificially prepared using a DNA synthesizer, or may be obtained by, for example, expressing a gene in the direction opposite to the usual case (*i.e.*, in the antisense direction). For details, see the descriptions in the section of the 21st embodiment of the present invention.

[0062] These antisense nucleotides are used for inhibiting the expression of Semaphorin Y as described hereinafter in the section of the 15th embodiment of the present invention, and are also useful as laboratory reagents for, for instance, *in situ* hybridization. In the present invention, "a chemically modified variant" specifically refers to such a variant that is chemically modified so as to enhance the transferability of the antisense nucleotide into cells or the stability of the antisense nucleotide in the cells. Examples of such chemically modified variant are phosphorothioate, phosphorodithioate, alkylphosphotriester, alkyl phosphonate, alkyl phosphoramidate and the like derivatives ("Antisense RNA and DNA", WILEY-LISS, 1992, pp. 1-50, *J. Med. Chem.*, **36**, 1923-1937 (1993)). The chemically modified variant may be prepared according to, for example, the references cited just above.

[0063] The 15th embodiment of the present invention is an antisense nucleotide, or chemically modified variant thereof, of the 14th embodiment described above, characterized in that it inhibits the expression of the protein of the 4th embodiment of the present invention.

[0064] mRNAs produced by usual gene transcription are sense-strands, and the antisense nucleotides or chemically modified variants thereof can bind to such sense-strand mRNAs in cells to inhibit the expression of those particular genes. Therefore, the above-described antisense nucleotides or chemically modified variants thereof can inhibit the expression of Semaphorin Y, and can thereby inhibit the activity of Semaphorin Y. Some of antisense nucleotides or chemically modified variants thereof having such effect will serve as CNS-neuron regeneration promoters as described hereinafter in the section of the 21st embodiment of the present invention.

[0065] It can easily be determined whether a particular antisense nucleotide prepared, or a chemically modified variant thereof, has a desired inhibitory effect or not, by directly introducing the antisense oligonucleotide itself or by introducing a gene which produces said antisense RNA when transcribed, into cells expressing Semaphorin Y, and then determining whether the amount of the expressed Semaphorin Y is decreased or not.

[0066] Examples of antisense nucleotide having such inhibitory effect are those oligonucleotides having sequences complementary to either the coding region or the 5' noncoding region of Semaphorin gene of the above-described embodiments. Especially preferred are those antisense nucleotides having sequences complementary to the transcription initiation site, translation initiation site, 5' noncoding region, exon-intron junction region, or 5' CAP region.

[0067] The 16th embodiment of the present invention is an antibody against the protein of the 4th or 6th embodiment, or against the peptide of any one of the 11th to 13th embodiments. Such antibody can easily be produced by using mouse or rabbit according to the procedures described in, for example, "Current Protocols in Immunology", pp. 2.4.1-2.6.6 (1992, J. E. Coligan ed.). Monoclonal antibodies can also easily be produced by the methods described in the above-mentioned reference. Such antibodies may be used in affinity chromatography or screening of cDNA libraries, and as pharmaceutical or diagnostic agents, or laboratory reagents. Some of such antibodies have the activity of neutralizing Semaphorin Y. Such neutralizing activity can easily be determined, as described above in the section of the 5th embodiment of the present invention, by adding Semaphorin Y to an activity assay system described in the section of the 1st embodiment of the present invention, and further adding thereto a test substance (*i.e.*, a candidate antibody against Semaphorin Y). Some of such neutralizing antibodies will serve as CNS-neuron regeneration promoters as described hereinafter in the section of the 21st embodiment of the present invention.

[0068] The 17th embodiment of the present invention is a pharmaceutical agent comprising, as an active ingredient, any one of all of the genes (DNAs), proteins, peptides, antisense nucleotides or chemically modified variants thereof, and antibodies of the present invention.

[0069] Among such pharmaceutical agents, CNS-neuron regenerators and neurite-outgrowth inhibitors for PNS-neuron will be described in the sections of the 21st and 22nd embodiments of the present invention, respectively. See, therefore, the sections of the 21st and 22nd embodiments for such applications.

[0070] It is being demonstrated in recent years that certain Semaphorins play important roles not only in the nervous system but also in non-nervous system. For example, it has been suggested that Semaphorin may probably act in inhib-

specific directions for use, dosage and the like, see the sections of the 21st and 22nd embodiments.

[0072] The 18th embodiment of the present invention is a method of screening for Semaphorin Y antagonists, char-

[0073] The screening is conducted by adding Semaphorin Y to an assay system for Semaphorin Y activity described in the prior art. The addition of Semaphorin Y to the assay system is in the presence of a test substance. The test substance is an antagonist, which inhibits, for example, the neurite outgrowth induced by Semaphorin Y. The addition of Semaphorin Y to the assay system is in the presence of a test substance. In particular,

the incubation period or only temporarily in the incubation period can be used as an indicator in the Semaphorin Y activity assay conducted with added Semaphorin Y. It is also important to confirm that the test substance alone does not inhibit the Semaphorin Y activity.

are fulfilled, one can consider the test substance as a Semaphorin-1 antagonist. However, to advance the test substance in the form of aqueous solution, an organic solvent such as DMSO may also be used as a solvent. In this case, it is important to minimize the volume of solvent so as to exclude any effects of the solvent on

and more preferably less than 1/100 volume relative to the culture medium. Some of Semaphorin Y antagonists thus obtained will serve as CNS-neuron regeneration promoters as described hereinafter in the section of the 21st embodiment.

of the 18th embodiment of the present invention. Such antagonist may have any structure and any form, provided that it inhibits the activity of Semaphorin Y.

prises the protein of the 6th embodiment, the peptide of any one of the 11th to 13th embodiments, or the antibody of the 16th embodiment of the present invention. In other words, it is a protein of the 6th embodiment, a polypeptide of any one of the 11th to 13th embodiments, or an antibody of the 16th embodiment of the present invention which has an

effect of inhibiting the activity of Serpiniform 1. Certain antagonists to the screening system of the 18th embodiment of the present invention, and some of the antagonists thus identified will serve as CNS-neuron regeneration promoters as described below in the section of the 21st embodiment of the

[0076] The 21st embodiment of the present invention is a CNS-neuron regeneration promoter, characterized in that it contains at least one of the antisense nucleotides or chemically modified variants thereof of the 14th or 15th embodiment of the present invention. Since this embodiment

relates to the use of substances in "regeneration therapy for CNS-neuron", specific effects and side effects of the substances are described below.

[0077] Application of antisense nucleotides has been attempted in various diseases, and in recent years, it is also intended to be applicable in neurological disorders (T/NS 20, No. 8, 321-322 (1997)).

nucleotide or chemically modified variant thereof of the 14th or 15th embodiment of the present invention can be used for inhibiting expression of Semaphorin Y gene. Accordingly, such antisense nucleotide may decrease the abundance of Semaphorin Y mRNA.

[0076] The present invention also provides a method for treating CNS diseases. Therapeutic methods using the nucleotide or

the variant include those in which the antisense oligonucleotide or its chemically modified variant is administered as such, and those in which antisense RNA is produced in cells.

especially preferably 12-25 bases. Antisense oligonucleotide or its chemically modified variant may be formulated by mixing it with stabilizing agent, buffer, solvent and the like prior to its administration. Such formulation may be co-administered with another therapeutic agent, such as an antineoplastic agent. Although the formulation thus prepared

disordered. Usually, regeneration of neuron takes several days to several months, and the formulation is administered

every day or every several days to several weeks during the period. To avoid such frequent administrations, a sustained-release mini-pellet formulation may be prepared and embedded near the affected site. Alternatively, a formulation may be gradually and continuously administered to the affected site by means of, for example, an osmotic pump. The dose is typically adjusted so that the concentration at the site of action will be 0.1 nM to 10 μ M.

[0080] In the method in which antisense RNA is produced in cells, a preferred antisense RNA has a length of, for example, more than 100 bases, preferably more than 300 bases, and more preferably 500 bases or more.

[0081] The methods by which a gene expressing an antisense RNA is introduced into a patient include an *in vivo* method in which the gene is directly introduced into cells in a living body, and an *ex vivo* method in which the gene is introduced into particular cells *ex vivo* and the cells are returned into the body (Nikkei Science, April, 1994, pp. 20-45; Gekkan-Yakuji, 36 (1), 23-48 (1994); Jikkenn-Igaku-Zokan, 12 (15), 1994; and references cited therein). An *in vivo* method is more preferred.

[0082] Such *in vivo* methods include a method employing recombinant viruses and other methods (Nikkei Science, April, 1994, pp. 20-45; Gekkan-Yakuji, 36 (1), 23-48 (1994); Jikken-Igaku-Zokan, 12 (15), in its entirety (1994); and references cited therein).

[0083] The methods employing recombinant viruses may include the methods in which Semaphorin gene is incorporated into a virus genome of, for example, retrovirus, adenovirus, adeno-associated virus, herpesvirus, vaccinia virus, poliovirus, or sindbis virus, and the recombinant virus is introduced into a living body. Among these methods, those employing retrovirus, adenovirus or adeno-associated virus are particularly preferred.

[0084] Other methods may include a liposome method or a lipofectin method. The liposome method is particularly preferred.

[0085] For the *ex vivo* methods, a micro-injection method, the calcium phosphate method, electroporation and the like may also be used, besides those techniques described above.

[0086] Administration of the gene to a patient is carried out via appropriate routes depending on particular disease or symptom to be treated, and the like. For example, it may be administered intravenously, intraarterially, subcutaneously, or intramuscularly, or directly administered into an affected site such as neuron. For example, when spinal cord is infected with the recombinant viruses, the expression of Semaphorin gene is inhibited exclusively in the spinal cord. Expression of antisense oligonucleotide of the present invention typically lasts several days to several months, and such single infection is sufficient to allow regeneration of neuron. The gene may also be reinfected, when weakly expressed. When administered by an *in vivo* method, the gene may be formulated in the form of, for example, a solution, and typically it is formulated in the form of an injection containing Semaphorin gene as an active ingredient to which conventional carrier and the like may be added, if necessary. In the case of liposomes or membrane-fused liposomes (such as Sendai virus (HVJ)-liposomes) containing Semaphorin gene, the liposome preparations may be in the form of a suspension, a frozen preparation, a centrifugally-concentrated frozen preparation or the like.

[0087] Although the amount of Semaphorin gene in the formulation may vary depending on the disease to be treated, the age and weight of the patient, and the like, it is typically 0.0001-100 mg, and preferably 0.001-10 mg, and such formulation is preferably administered once every several days to several months.

2) Modified protein of Semaphorin Y

[0088] As described above in the sections of the 5th and 6th embodiments of the present invention, one can prepare a modified Semaphorin Y of which neurite-outgrowth inhibition activity on CNS-neuron has been eliminated. When administered into a living body, such modified protein is expected to bind to receptors for Semaphorin Y in place of Semaphorin Y, resulting in inhibition of Semaphorin Y activity and promotion of regeneration of CNS-neuron.

[0089] Such modified protein of Semaphorin Y is formulated with stabilizer, buffer, and diluent, and administered to a patient for therapy. Such formulation may be administered via various routes, and it is preferred to topically administer to the focal site. Since regeneration of neuron typically takes several days to several months, the formulation is administered once or more in order to continuously inhibit Semaphorin Y activity throughout the period. When administered more than once, it is desirable to administer it every day or repeatedly at appropriate intervals. When administered to CNS by injection, for example, into spinal cord, several hundreds μ g to 2 g, preferably less than several tens mg, are used per administration. To reduce the administration frequency, it may be administered using a sustained-release formulation or gradually administered over a long period by means of, for example, an osmotic pump. Alternatively, it may be administered by grafting cells expressing such modified Semaphorin Y protein into a living body.

3) Peptide of Semaphorin Y

[0090] Some of the peptides of any one of the embodiments from 11th to 13th of the present invention suppress the neurite outgrowth inhibition activity of Semaphorin Y on CNS-neuron by inhibiting the binding of Semaphorin Y to its receptors, resulting in promotion of CNS-neuron regeneration. Examples of peptide having such effect include a pep-

tide characterized in that it contains aspartic acid residue at position 198 of human Semaphorin Y shown in SEQ ID NO: 6 or an amino acid residue at the position corresponding to that of said aspartic acid residue, as described above in the section of the 13th embodiment of the present invention. The suppression may be any one of competitive, noncompetitive, uncompetitive, and allosteric inhibitions.

- 5 [0091] As for the methods of formulating or administering such polypeptides, and their doses, see the above section "2) Modified protein of Semaphorin Y".

4) Antibody against Semaphorin Y

- 10 [0092] A neutralizing antibody which neutralizes the activity of Semaphorin Y is expected to promote the regeneration therapy of CNS-neuron by inhibiting Semaphorin Y activity, when administered into a living body.

[0093] The methods of formulating or administering such neutralizing antibodies and their doses may be the same as described in the above section "2) Modified protein of Semaphorin Y". Alternatively, a method in which cells producing a monoclonal antibody are grafted directly into CNS may also be used, as described in *Nature*, 343, 269-272 (1990).

- 15 [0094] The 22nd embodiment of the present invention is a neurite outgrowth inhibitor for PNS-neuron, characterized in that it contains at least one of the proteins of the 4th embodiment of the present invention. Although the proteins of the 4th embodiment of the present invention may inhibit the neurite outgrowth of CNS-neuron, they are also expected to inhibit the neurite outgrowth of PNS-neuron, since PNS-neuron also probably expresses receptors for Semaphorin Y, and receptors for other Semaphorins also probably react with Semaphorin Y. Accordingly, they may serve as therapeutic agents for atopic dermatitis, pain or other diseases by virtue of their inhibition activity on neurite outgrowth of PNS-neuron.

20 [0095] As for the methods of formulating or administering such proteins, and their doses, see the above section "2) Modified protein of Semaphorin Y".

- 25 [0096] The 23rd embodiment of the present invention is a transgenic animal in which either the gene of any one of the 1st to 3rd and 5th embodiments, or DNA of the 7th embodiment of the present invention has been artificially inserted into its chromosome, or has been knocked out.

[0097] As apparent from the following references, one skilled in the art can quite easily produce a transgenic animal which expresses the gene of the 1st, 4th, 7th, or 9th embodiment of the present invention, in the light of the gene information on Semaphorin Y of the present invention: "Manipulation of Mouse Embryo" edited by B. Hogan *et al.*, 1986, Cold Spring Harbor Laboratory; Shinichi Aizawa, "Gene Targeting", 1995, Yodosha, etc. Accordingly, the transgenic animal thus produced is naturally included within the scope of the present invention. The transgenic animal thus produced is very useful as an animal model for developing pharmaceuticals or as an animal used for screening of pharmaceuticals. Furthermore, a so-called knockout animal in which the gene of the 1st, 4th, 7th, or 9th embodiment of the present invention has been deleted is characterized in that it does not contain such gene. As described in literatures, or as

35 apparent from the common knowledge in the art, such knockout animals cannot be produced without the gene information on Semaphorin Y of the present invention. It goes without saying, therefore, that such knockout animals are included within the scope of the present invention.

- [0098] While Semaphorin Y has an important *in vivo* function relating to regeneration of neurons as described above, it has been also suggested as mentioned above that Semaphorin Y may have other unknown functions such as immunosuppression (*Cell*, 75, 1389-1399 (1993)). Accordingly, it is quite important to investigate the expression of Semaphorin Y gene or the distribution and function of Semaphorin Y protein for studying this technical field or for diagnosing patients with neurological disorders or other diseases. The present invention can also provide gene probes, antibodies, recombinant proteins, transgenic animals and the like which can be used for such purposes.

45 BRIEF DESCRIPTION OF DRAWINGS

[0099] Fig. 1 shows a picture of electrophoresis indicating distribution of Semaphorin Y expression among various tissues determined by Northern analysis.

- 50 [0100] Total RNAs were extracted from various tissues of six-weeks old rats, electrophoresed on 1% agarose-formamide gel, blotted onto a filter, and hybridized with a ³²P-labeled rat Semaphorin Y DNA probe to determine the distribution of Semaphorin Y mRNA expression. Fifteen µg of RNA was loaded in each lane. The upper panel shows the result of autoradiography. The positions corresponding to 18S and 28S ribosomal RNAs are indicated at the left margin of the panel. The lower panel shows the ethidium bromide staining of the gel. The upper and lower bands correspond 28 and 18S ribosomal RNAs, respectively.

- 55 [0101] Fig. 2 shows a picture of electrophoresis indicating distribution of Semaphorin Y expression among CNS tissues determined by Northern analysis.

[0102] Total RNAs were extracted from CNS tissues of six-weeks old rats, electrophoresed on 1% agarose-formamide gel, blotted onto a filter, and hybridized with a ³²P-labeled rat Semaphorin Y DNA probe to determine the distribution of

Semaphorin Y mRNA expression. Fifteen μ g of RNA was loaded in each lane. The upper panel shows the result of autoradiography. The positions corresponding to 18S and 28S ribosomal RNAs are indicated at the left margin of the panel. The lower panel shows the ethidium bromide staining of the gel. The upper and lower bands correspond 28 and 18S ribosomal RNAs, respectively.

[0103] Fig. 3 shows a picture of electrophoresis indicating distribution of Semaphorin Y mRNA expression among human CNS tissues determined by Northern analysis.

[0104] A Membrane filter onto which mRNAs prepared from various regions of human CNS tissues have been transferred after being electrophoresed (2 μ g/lane) (Clontech) was hybridized with 32 P-labeled Semaphorin Y DNA probe to determine the distribution of Semaphorin Y mRNA expression. The figure shows the result of autoradiography. In this figure, the arrows indicate the positions of Semaphorin Y mRNA bands. Positions of size makers are indicated in kb at the left margin of the figure.

[0105] Fig. 4 shows a picture of electrophoresis indicating expression of Semaphorin Y protein in COS 7 cells.

[0106] An expression plasmid for Semaphorin Y having additional 10 amino acids derived from human c-Myc added at its C-terminus was constructed, and introduced into COS 7 cells for transient expression (indicated as rSYmyc). A plasmid containing no Semaphorin Y gene was used as control (indicated as Control). At day 3 after introducing plasmids, the cells were harvested, and the membrane fraction was prepared. The membrane fraction was fractionated by SDS-PAGE, and then subjected to Western blotting using an anti-Myc antibody. In this figure, the arrow indicates the position of the band of Semaphorin Y protein having added Myc peptide. Positions and molecular weights of size markers are indicated in kD at the left margin of the figure.

[0107] Fig. 5 shows a picture of electrophoresis indicating the *in vivo* distribution of Semaphorin III expression among various tissues determined by Northern analysis.

[0108] Total RNAs were extracted from various tissues of adult rats, electrophoresed on 1% agarose-formamide gel, blotted onto a filter, and hybridized with 32 P-labeled mouse Semaphorin III DNA probe to determine the distribution of Semaphorin III mRNA expression. Fifteen μ g of RNA was loaded in each lane. The upper panel shows the result of autoradiography. The positions of 18S and 28S ribosomal RNAs are indicated at the left margin of the figure. The lower panel indicates the ethidium bromide staining of the gel. The upper and lower bands correspond to 28S and 18S ribosomal RNAs, respectively.

EXAMPLES

[0109] Fundamental procedures for experiments are described in detail in many publications such as "Molecular Cloning, 2nd Ed." edited by Maniatis *et al.* (Cold Spring Harbor Laboratory Press, 1989), "Current Protocols in Molecular Biology" edited by Ausubel *et al.* (John Wiley & Sons, 1987), and "Saibo-Kogaku-Jikken Protocols" edited by Department of Oncology, The Institute of Medical Science, The University of Tokyo (Shujunsha, 1991). The present invention is not intended to be limited by the following examples, and the examples may be of course modified as usual.

Example 1

Isolation of rat Semaphorin Y gene

(1) Search through database for a novel Semaphorin gene

[0110] Using the dbEST database of the National Center for Biotechnology Research (Bethesda, MD, US), search was performed for a sequence which encodes an amino acid sequence relatively well conserved in known Semaphorin genes and which is found in only cDNAs from postnatal brain but not in cDNAs from peripheral tissues. As a result, the base sequence of File No. R59527 proved to encode a sequence consisting of seven amino acids common to known Semaphorin genes (Gln (or Arg)-Asp-Pro-Tyr-Cys-Ala (or Gly)-Trp). However, the sequence information of R59527 consisting of 238 bases is so short compared with the cDNAs for known Semaphorin genes, and only several percent of the total bases could be translated to a sequence common to those in known Semaphorins. In addition, the reading frame could not be determined because the sequence of R59527 is not the one finally determined. It was, therefore, impossible to conclude that the base sequence of R59527 is part of a novel Semaphorin gene. Then, the inventors firstly confirmed that a gene containing the above sequence is expressed in the adult brain, and then sought to clone the full length cDNA containing the above sequence and determine its gene structure.

(2) Confirmation of the expression of the gene containing the sequence of R59527 in the brain

[0111] To confirm that the gene is expressed in the adult human CNS, two DNA primers bounding a segment of about 170 bp, 5' TGGCTGTATTGTCTACCT 3' (SEQ ID NO: 8) and 5' TGGATTCCTGGTTCNAGCC 3' (SEQ ID NO: 9), were

synthesized on the basis of the base sequence of R59527, and used in PCR under conventional conditions together with cDNAs prepared from a human brain cDNA library (Clontech) as templates. As a result, about 170 bp fragment was amplified as expected. The DNA was then cloned into pCRII (Invitrogen) according to the manufacturer's protocol, and the base sequence was determined to confirm that the fragment has the same base sequence as that of R59527. More than 98% of the sequence thus obtained (SEQ ID NO: 7) coincided with that of R59527, confirming that a gene containing the sequence of R59527 is expressed in the adult human brain.

(3) Isolation of rat Semaphorin Y gene

[0112] Using the 170 bp fragment cloned in (2), which corresponds to part of R59527, as a probe, the inventors cloned a full-length cDNA containing the sequence of the probe and determined the structure. Since preparation of rat cDNA library is easier than that of human cDNA library, the rat gene was firstly cloned. A cDNA library was prepared by conventional methods described in the above-mentioned laboratory manuals, using mRNAs prepared from rat brain and muscle by conventional procedures with Lambda Zap II (λ ZapII) cDNA Library Preparation Kit (Stratagene). About 150 thousand plaques were then generated on agar plates using the cDNA library, and the plaques were transferred onto nylon membranes (Nippon Pall). After denaturing and neutralizing the DNAs, they were fixed with ultraviolet rays of 0.6 J/cm², and used in hybridization. The hybridization was conducted by placing the nylon membrane and the 170 bp DNA fragment labeled with ³²P (prepared using Megaprime DNA Labeling System (Amersham)) as a probe in a hybridization buffer (45% (v/v) formamide, 5x SSPE (1x SSPE consists of 0.15 M sodium chloride, 10 mM sodium dihydrogenphosphate, and 1 mM ethylenediaminetetraacetic acid disodium salt, adjusted to pH 7.0), 2x Denhardt's solution (Wako Pure Chemical Industries), 0.5% (w/v) sodium dodecyl sulfate (SDS), 20 μ g/ml salmon sperm DNA (Wako Pure Chemical Industries)) and allowing them to stand at 42°C for 48 hours. After the reaction, the nylon membrane was washed 2-3 times in 2x SSPE, 0.5% (w/v) SDS at room temperature for 10 min, and then 2-3 times in 2x SSPE, 0.5% (w/v) SDS at 42°C for 10 min. The filters thus prepared were analyzed using BAS 2000 Bio-Imaging Analyzer (Fuji Film), and 6 positive signals were obtained. Plaques corresponding to the positive signals were excised from the agar plates, placed in 500 μ l of SM buffer (100 mM sodium chloride, 15 mM magnesium sulfate, 50 mM Tris (pH 7.5), 0.01% gelatin) supplemented with 20 μ l of chloroform, and left stand overnight at 4°C to elute the phages. The recombinant lambda phages thus obtained were subjected to a secondary screening according to the procedures as described above, and single plaques were isolated. The phages thus obtained were treated in the following manner for *in vivo* excision of a phagemid containing the cDNA insert, according to the protocols supplied by Stratagene. Agar gels containing the 4 single plaques obtained in the secondary screening were each placed in 500 μ l of SM buffer, supplemented with 20 μ l of chloroform, and then allowed to stand overnight at 4°C. Two hundred fifty μ l of the phage solution obtained, 200 μ l of *E. coli* XL-1 Blue MRF' suspended in 10 mM magnesium chloride at OD₆₀₀ = 1.0, and 1 μ l of ExAssist helper phage (>1x10⁶ pfu/ml) were mixed, and incubated at 37°C for 15 min. Then, 3 ml of LB medium (prepared by mixing 0.5% (w/v) sodium chloride, 1% (w/v) Bactotrypton (Difco), and 0.5% (w/v) yeast extract (Difco) and the mixture was adjusting to pH 7.0 using 5 M sodium hydroxide) was added, and the mixture was shaken at 37°C for 2-3 hours. The cells were removed by centrifuging at 2000xg for 15 min, and the supernatant was heat-treated at 70°C for 15 min. The supernatant was then centrifuged again at 2000xg for 15 min, and recovered as a stock solution of a phagemid containing the cDNA insert. An aliquot (10-100 μ l) of the phagemid stock solution was mixed with 200 μ l of *E. coli* SOLR (OD₆₀₀ = 1.0), incubated at 37°C for 15 min. Then, 10-50 μ l of the mixture was plated onto an ampicillin plate, and incubated overnight at 37°C to obtain *E. coli* strain which contained the phagemid corresponding to the above positive plaque.

(4) DNA sequencing

[0113] The base sequence of the cDNA clone thus obtained was analyzed on Perkin-Elmer Model 377 DNA Sequencer to determine the complete base sequence. The reaction was carried out using PRISM Dye termination kit (Perkin-Elmer). The DNA base sequence thus determined (3195 bases), the putative open reading frame (2787 bases), and the amino acid sequence (929 amino acids) are shown in SEQ ID NOs: 1, 2, and 3, respectively. [0114] The protein contained a so-called semaphorin domain at positions 46 through 570 in the amino acid sequence, definitely confirming that the protein belongs to the Semaphorin family. The protein encoded by the gene was thus designated Semaphorin Y. In addition, since the base sequence of positions 1574 through 1811 in the Semaphorin Y gene shown in SEQ ID NO: 1 had 89% identity with the whole sequence of R59527 consisting of 238 bp, it was confirmed that R59527 is a partial sequence of human Semaphorin Y gene.

Example 2Distribution of Rat Semaphorin Y expression determined by Northern analysis

5 [0115] In order to determine the distribution of Semaphorin Y gene expression among rat tissues, RNAs were prepared from various tissues and used in Northern analysis. RNAs were prepared as follows using various rat tissues according to AGPC method (Takashi Tuji and Toshikazu Nakamura, *Jikken-Igaku*, vol. 9, 1991, pp. 1937-1940; M. F. Ausubel *et al.* ed., "Current Protocols in Molecular Biology", 1989, pp. 4.2.4-4.2.8, Greene Pub. Associates & Wiley-Interscience). Briefly, 10 ml of a denaturing solution (4M guanidine thiocyanate, 25 mM sodium citrate (pH 7.0), 0.5% Sarkosyl, 0.1 M 2-mercaptoethanol) was added to each 1 g of excised tissues, and quickly homogenized using a Polytron homogenizer. To the homogenate, 0.1 volume of 2 M sodium acetate (pH 4.0), 1 volume of water-saturated phenol, and 0.2 volumes of chloroform-isoamyl alcohol (49:1) were added, and the mixture was vigorously stirred. After centrifugation, the aqueous layer was isolated, an equal volume of isopropyl alcohol was added thereto, and the mixture was allowed to stand at -20°C for 1 hour. The precipitate was recovered by centrifugation, and dissolved again in 2-3 ml of the denaturing solution per 1 g tissue. An equal volume of isopropyl alcohol was added, and the mixture was allowed to stand at -20°C for 1 hour, and then RNA was centrifuged. The precipitate was washed with 75% ethyl alcohol, dried briefly, and then dissolved in an appropriate amount of water.

[0116] Subsequently, electrophoresis and Northern blotting of RNAs were performed by conventional methods described below. RNAs prepared from various tissues were firstly electrophoresed on 1% agarose gel containing formaldehyde. The gel was shaken in 50 mM NaOH for 20 min, and then in 10x SSPE for 40 min. The RNAs were then blotted onto a nylon membrane (Biodyne B, Nippon Pall) by means of capillary transfer, and fixed using a UV cross-linker (Stratagene) (0.6J/cm²) for use in hybridization. A probe was prepared as follows. Firstly, PCR was carried out using two primers, 5' TGTGTAAACGTGACATGG 3' (SEQ ID NO: 10) and 5' TGCTAGTCAGAGTGAGGA 3' (SEQ ID NO: 11), with rat Semaphorin Y cDNA obtained in Example 1 as template to amplify a fragment of 477 bp. This fragment was cloned into pCR II in the same manner as described above, and the base sequence was determined to confirm that it was a fragment of rat Semaphorin Y gene. Using this plasmid DNA as template, PCR was carried out in conventional manner with the above primers to amplify the aimed fragment of 545 bp. The amplified DNA separated and purified using agarose gel was labeled with ³²P using Megaprime DNA Labeling System (Amersham) as described in Example 1, and used as a probe. Hybridization was carried out by placing the nylon membrane and the probe DNA in the same hybridization buffer as described above in (2) and allowing them to stand at 42°C for 48 hours. After the reaction, the nylon membrane was washed 2-3 times in 2x SSPE, 0.5% (w/v) SDS for 10 min at 42°C, and then 2-3 times in 2x SSPE, 0.5% SDS (w/v) at 55°C for 10 min. Radioactivity on the membrane was then analyzed using BAS 2000 Bio-Imaging Analyzer. As shown in Figs. 1 and 2, the result demonstrated that mRNA for Semaphorin Y was widely expressed in the adult CNS, whereas the expression was not detected in peripheral tissues with the only exception of muscle, exhibiting the characteristic features expected with Semaphorin gene as CNS-neuron regeneration inhibitor.

Example 3Sequence determination of human Semaphorin Y

40 [0117] Since R59527 has proved to be part of human Semaphorin Y gene as described above, an EST clone containing the sequence of R59527 (#41581) was obtained from Genome Systems Inc. (US), and the complete base sequence was determined by the method described above. The determined base sequence had a high homology to the entire base sequence for rat Semaphorin Y shown in SEQ ID NO: 1 with 74% of the bases being the same. In addition, the 5' region of the base sequence contains a stretch, presumably part of the open reading frame, which could be continuously translated into 427 amino acids. This amino acid sequence had an identity of 82% with that of the region from position 504 to position 929 of rat Semaphorin Y gene shown in SEQ ID NO: 3, indicating that the sequence was certainly part of human Semaphorin Y. However, the sequence corresponding to the N-terminal of human Semaphorin Y could not be determined from this clone #41581. In order to determine the base sequence for human Semaphorin Y in full length, human hippocampus and forebrain cDNA libraries purchased from Stratagene were screened as described above using various rat Semaphorin Y cDNA fragments as probes to obtain a clone #10. The base sequence of the clone #10 determined by the same procedures as described above overlapped with the above clone #41581 by about 200 bases, and further contained in its 5' region a cDNA sequence consisting of more than 1700 bases. The complete base sequence (3432 bases) constructed from #41581 and #10, the open reading frame (2790 bases), and the amino acid sequence (930 amino acids) for human Semaphorin Y are shown in SEQ ID NOs: 4, 5, and 6, respectively.

55 [0118] *E. coli* strain SOLR (hSY10), a transformant obtained by introducing the plasmid hSY10, which incorporates the insert of the above clone #10 (the region corresponding to cDNA for human Semaphorin Y) in a vector pBluescript,

into *E. coli* strain SOLR, has been deposited at The National Institute of Bioscience and Human Technology (1-1-3 Higashi, Tsukuba, Ibaraki, Japan) under Deposit No. FERM BP-6021 on July 11, 1997.

- [0119] *E. coli* strain DH10B (N041581), a transformant obtained by introducing the plasmid N041581, which incorporates the insert of the above clone #41581 (the region corresponding to cDNA for human Semaphorin Y) in a vector Lafrmid BA, into *E. coli* strain DH10B, has been deposited at The National Institute of Bioscience and Human Technology (1-1-3 Higashi, Tsukuba, Ibaraki, Japan) under Deposit No. FERM BP-6022 on July 11, 1997.

Example 4

10 Distribution of human Semaphorin Y expression determined by Northern analysis

[0120] Northern analysis was performed as described in Example 2 with human mRNA blotting membrane (Clontech) using a rat Semaphorin Y cDNA fragment consisting of 479 bp from position 832 to position 1310 in SEQ ID NO: 1 obtained by PCR as a probe to determine the distribution of Semaphorin Y mRNA expression among various regions

15 in human adult CNS tissues. As shown in Fig. 3, human Semaphorin Y mRNA was widely expressed in various regions of the adult CNS tissues, and a particularly high expression was observed in the cerebellum.

[0121] As stated above, Semaphorin Y is widely expressed in human CNS tissues as is the case with its rat homologue, indicating that Semaphorin Y may be responsible for functions common to rodents and primates.

20 Example 5

Expression of Semaphorin Y in animal cells

[0122] A fragment encoding Myc tag having the sequence Asp-Ile-Gly-Gly-Glu-Gln-Lys-Lue-Ile-Ser-Glu-Glu-Asp-Leu was inserted just before the stop codon of rat Semaphorin Y gene, and the recombinant gene was introduced into an expression plasmid pUCSR α . The expression plasmid was transfected into COS 7 cells according to DEAE-dextran method ("Current Protocols in Molecular Biology" edited by F. M. Ausubel, John Wiley & Sons, 1987), and the cells were harvested with a cell scraper after 48 hours. Harvested cells were homogenized in the presence of Solution A containing protease inhibitors (Hank's physiological saline containing 10 mM HEPES pH 7.4, 1 mM EDTA, 50 μ M leupeptin, 2

30 μ M pepstatin, 0.5 mM PMSF, and 7.8 mTIU/ml aprotinin), and the homogenate was separated into precipitate and supernatant by high-speed centrifugation at 12,000g for 10 min. The precipitate from the high-speed centrifugation which contained the membrane fraction was washed twice with Solution A, suspended in 2 volumes of 2.25 M sucrose/PBS, and overlaid onto 2.25 M sucrose/PBS. After 0.8 M sucrose/PBS was further overlaid onto the top, it was centrifuged at 12,000g for 20 min. The membrane fraction was recovered from the lower interface, further washed

35 twice, and stored at -80°C until use.

[0123] The membrane fraction obtained was subjected to SDS-PAGE (10%-20% gradient gel), and then to Western blotting in conventional manner to confirm the production of Semaphorin Y of the present invention. During this procedure, an anti-Myc antibody 9E10 (Calbiochem) and an alkaline phosphatase-labeled anti-mouse IgG antibody (Biosource) were used as the primary and secondary antibodies, respectively. The result of Western blotting showed a

40 specific band at the position corresponding to about 130 kDa as shown in Fig. 4, confirming that Myc-tagged rat Semaphorin Y protein was expressed in COS cells and existed in the membrane.

Example 6

45 Activity measurement of Semaphorin Y

[0124] The membrane fraction obtained in Example 5 and another membrane fraction prepared in the same manner from COS 7 cells untransfected with Semaphorin Y are each added to culture medium for neurons such as CNS-neurons or dorsal root ganglion cells, and the growth-cone collapse activities are compared by the method described in M. Igarashi *et. al.*, Science, 259, 77-79 (1993). The result demonstrates that the membrane fraction from COS 7 cells

50 transfected with the expression plasmid for Semaphorin Y has a significantly high growth-cone collapse activity.

Reference example 1

55 [0125] Identification of the site essential to the Semaphorin activity using Semaphorin III

[0126] PCR was conducted on the basis of the sequence information on Semaphorin III described in Neuron, 14, 941-948 (1995), and the structural gene of Semaphorin III was incorporated into an expression plasmid pUCSR α . The expression plasmid was then introduced into COS 7 cells by DEAE-dextran method. After 2 days, the Semaphorin III

activity contained in the culture supernatant was determined by a method similar to that described in *Cell*, 75, 217-227 (1993), using the growth-cone collapse activity on chicken dorsal root ganglion cells as an indicator. As a result, one clone which did not exhibit any activity was found. The base sequencing of the clone revealed that aspartic acid residue at position 198 was substituted by glycine. When compared with other known animal Semaphorins, the regions before and after the position 198 were not markedly conserved, although the position corresponding to aspartic acid was highly conserved among Semaphorins with a few exceptions in which glutamic acid was located at that position. This suggested that the aspartic acid residue is essential to expression of the activity. The gene was then subjected to a site-directed mutagenesis by a conventional method to replace the glycine residue with aspartic acid. Since this mutagenesis restored the strong collapse activity, it was confirmed that all of the regions in the expression plasmid normally function except for that position. In conclusion, the aspartic acid at position 198 of Semaphorin III appears essential to expression of the Semaphorin function. The amino acid residues corresponding to the aspartic acid are aspartic acid at position 197 in the amino acid sequence of rat Semaphorin Y shown in SEQ ID NO: 3, and aspartic acid at position 198 in the amino acid sequence of human Semaphorin Y shown in SEQ ID NO: 6.

Reference example 2

Tissue-specific gene expression of Semaphorin III determined by Northern analysis

[0127] To determine the distribution of Semaphorin III gene expression among mouse tissues, RNAs were prepared from various adult mouse tissues, and subjected to Northern analysis. The procedures for preparation, blotting, and hybridization of RNA were the same as those described in Example 2. As a probe, the 560 bp *Msp*I fragment of mouse Semaphorin III DNA described in Reference example 1 was used. As a result, it was demonstrated as shown in Fig. 5 that the expression of Semaphorin III in the adult is extremely high in the lung, while it is rather low in the CNS.

EFFECTS OF THE INVENTION

[0128] The present invention provides Semaphorin Y inhibiting neurite outgrowth, and a gene therefor, as well as other Semaphorins hybridizing to said Semaphorin Y gene, modified proteins or partial peptides of said Semaphorin Y, antibodies against said Semaphorin Y, antisense nucleotides against said Semaphorin Y gene, and the use of such substances as pharmaceutical or diagnostic agents or laboratory reagents. The present invention further provides a method of screening for Semaphorin Y antagonists employing said Semaphorin Y, Semaphorin Y antagonists obtained by said screening method, pharmaceutical agents comprising such antagonists, and transgenic animals involving said Semaphorin Y.

SEQUENCE LISTING

5 SEQ ID NO: 1

SEQUENCE LENGTH: 3195 base pairs

SEQUENCE TYPE: nucleic acid

10 STRANDEDNESS: double

TOPOLOGY: linear

15 MOLECULE TYPE: cDNA to mRNA

HYPOTHETICAL: No

ANTI-SENSE: No

20 ORIGINAL SOURCE:

ORGANISM: rat (*Rattus norvegicus*)

25 TISSUE TYPE: brain

FEATURE:

30 FEATURE KEY: 5' UTR

LOCATION: 1..50

IDENTIFICATION METHOD: E

35

FEATURE KEY: CDS

40 LOCATION: 51..2837

IDENTIFICATION METHOD: E

45

FEATURE KEY: 3' UTR

LOCATION: 2838..3195

50 IDENTIFICATION METHOD: E

SEQUENCE DESCRIPTION:

55

GCGGCCCGGT CGACGGGGTA CTCCTGGGG CTTGAGCTGC CCGGCACAGG ATGCCCGGTG 60
 CCCCCACTC CATGCCCTTG CTGCTGCTGT TGCTGCTGT ACTCCCCAA GCCCAGACTG 120
 CCTTTCCCA GGACCCATC CCTCTGTGA CCTCTGACCT ACAAGGTACC TCTCCGTCAT 180
 CCTGGTTCCG GGGCCTGGAG GACGATGCTG TGGCTGCGGA ACTTGGGCTG GACTTTCAGA 240
 GATTCTGAC CTGAACCGG ACCTTGCTTG TGGCTGCCCG GGATCACGTT TTCTCCTTCG 300
 ATCTTCAAGC CCAAGAAGAA GGGGAGGGGC TGGTGCCAA CAAGTTTCTG ACATGGCGGA 360
 GCCAAGACAT GGAGAATTGT GCTGTCCGG GAAAGCTGAC GGACGAATGC TACAACTACA 420
 TCCGTGTTCT TGTCCCTGG GACTGCGAGA CACTCCTTG CTGTGAACA AATTCCTTCA 480
 GCCTGTGTG TCGCAGCTAT GGGATAACAT CTCTGCAACA GGAGGGTGAG GAGCTGAGTG 540
 GGCAAGCTCG ATGCCCTTT GATGCCACC AGTCCACTGT GGCCATCTCT GCAGAGGGTA 600
 GTTTGTACTC AGCCACAGCA GCAGATTTC AGCCAGTGA TGCTGTGTT TACAGAAGCC 660
 TTGGACCTCA CCCCCACTC CGTTCTGCAA AGTATGACTC CAAGTGGCTT CGAGAGCCAC 720
 ACTTTGTCTA TGCTTTGGAG CATGGAGACC ATGTCTACTT CTTTCTTCG GAGAAGTCTC 780
 TGTGGAGGAC GCCCGGCTG GGGAGGGTGC AGTTTTCCG GGTGGCCCG GTGTGTAAAC 840
 GTGACATGGG TGGCTACCA CGGGCCTTG ATCGCCACTG GACATCCTC CTTAAGCTGA 900
 GGTCAACTG CTCCGTCCCT GGGGACTCTA CTTTCTACTT TGATGTCTTA CAGTCCTTAA 960
 CTGGGCTGT GAACCTGCAT GGGGCTCTG CCTCTTTGG GGTCTTCACT ACTCAGACCA 1020
 ATAGCATTCC TGGGTCTGCA GTCTGCGCT TCTACCTAGA TGACATTGAA CGTGGCTTTG 1080
 AGGGCAAGTT CAAGGAGCAG AGGAGTCTGG ATGGGCTG GACTCCTGTG TCTGAGGACA 1140
 AAGTCCCTC ACCCAGGCCA GGTCTCTGTG CAGGTGTGG TGCAGTGCC TTATTCTCCT 1200
 CCTCTAAGA CCTGCTGAC GATGTCTGC TCTTCATCA GGCACACCA CTGCTGGATC 1260
 CCGCTGTGCC ACCTGCCACC CATCAACCTC TCCTCACTCT GACTAGCAGG GCTCTACTGA 1320
 CCCAGGTAGC TGTGATGGT ATGGCTGGC CCCACAGAA TACTACAGTC CTGTTTCTTG 1380
 GCTCCAATGA TGGACAGTG CTGAAGGTGC TACCTCCAG GGCACAGTCT CTGGGACCG 1440
 AGCCTATCAT ATTGAAGAG ATTGATGCCT ACAGCCATG CCGGTGCAGT GGAAGCGGT 1500

CACCCCGAGC TGCTCGACGG ATCATAGGGC TGGAGCTGGA CACTGAGGGT CACAGGCTTT 1560
 TTGTGGCCTT TCCTGGATGC ATCGTCTACC TCTCTCTCAG CCGCTGTGCC CGGCATGGAG 1620
 5 CATGTCAGAG GAGCTGCCTG GCTTCTCTGG ACCCATACTG TGGATGGCAT CGGTTCCGAG 1680
 GCTGTGTGAA TATCAGGGGA CCTGGAGGGA CTGATGTGGA TCTGACTGGG AACCAGGAAT 1740
 10 CCATGGAGCA TGCTGACTGC CAAGATGGAG CGACTGGGAG TCAGTCTGGC CCTGGAGATT 1800
 CTGCCATATGG CGTCCGAGG GACCTTTCCC CAGCCTCAGC CTCCCGATCC ATCCCATCC 1860
 CACTCCTCCT GGCCTGTGTG GCGGCGGCCT TCGCTTTGGG CGCCTCAGTC TCCGGCCTCT 1920
 15 TGGTGTCTG TGCTTGTCT CGCGGAACC GCCGTCCGAG CAAGGACATC GAGACCCCG 1980
 GGCTGCCGCG CCCCCTCTCC CTTCGCAGTC TGGCGAGGCT GCACGGTGGC GGTCTGAGC 2040
 20 CCCCCTCTCC GCGCAAGGAT GGTGATGCAG CGCAAACGCC CCAGCTCTAC ACTACCTTCC 2100
 TGCTCCGCC CGAGGGCGGA TCCCACCGG AGCTGGCCTG CCTGCCACC CCGGAGACCA 2160
 CGCCGAGCT GCCGTGAAG CACCTCCGTG CCTCCGGGG TCCCTGGGAG TGAACCAGA 2220
 25 ACGGAACAA CGCTTCGGAG GCGCCAGGCC GCGCACGGG CTGCAGCGCG GCGGGCGGGC 2280
 CCGCCCCGCG CGTGTGGTG AGGCCACGC CCCCTGGCTG CCGGGGAG GAGGTGGAGG 2340
 30 TGACCACGCT GGAGAACTG CTGCGCTACC TGCACGGCCC GCAGCGGCC AGGAAGGCA 2400
 GCGAACCTCT CGCCTCCGC CCGTTCACT CCGGCCGCC TGCTCGGAG CCGGCGCGC 2460
 35 CTTGTTCGT GGACTCCAGC CCGATGCCTC GTGATTGCT GCGGCGCTG AGGCTCGAG 2520
 TACGCCCCGA CGGCAAGGC GCGCCCCGA GCGGCGGCC TGCTCTCTG GCGCCGGCTC 2580
 CACGCTGGG CGTCAGGCG AGCGAAGAT TGCCCTTCCC CACGCACCG GCGCCCCCG 2640
 40 GCCTGCTCAC CCGAGTCCC TCGGAGGCC CGTCCAGTA CTCGGGGGG CCGGGAGGC 2700
 ACCTCCTGTA CCTGGGCCG CCGACGCC ACCGCGCCG CTCCTGAAG AGGTGGAGC 2760
 45 TGAAGTCTC ACTGTGCCC AAACGCCCC TCGCCACACC GCGCAGCCC GCGCGCACG 2820
 GCAGCCATTT TAATTCTGA CAGAAGCTG TAGCGCCGT CGAGGCGCTG GAGGCTAGG 2880
 50 CCTGCGAGG CCGCTGGCT TCCCGACTC CAAGACTCT CCGGGTCCC CTCTCGCCTC 2940
 GTTTATTTA TTACTGTCT TCCCCTGT CTTTGGCGA GGAGCTGCC GCTCGAGCG 3000

CCAGCATTTC AGGGGACCTG GCCGACTCCC ACTCCCCGCT CCTTCCAGC CACGCTGCCT 3060
TAACTCGTOG CTCGGGACTC CCGCGGACTG GGGCCCGGGC GGGCCGGCCG GGGCTGGAGC 3120
CGCGCGCTGT GTACAGAGTC CTCGGGCTC CTGGGGCCGG GACGTGCCTC CTCCTACTGT 3180
GTAGGAGCCC CCACC 3195

SEQ ID NO: 2

SEQUENCE LENGTH: 2787 base pairs

SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: cDNA to mRNA

HYPOTHETICAL: No

ANTI-SENSE: No

ORIGINAL SOURCE:

ORGANISM: rat (*Rattus norvegicus*)

TISSUE TYPE: brain

FEATURE:

FEATURE KEY: CDS

LOCATION: 1..2787

IDENTIFICATION METHOD: E

SEQUENCE DESCRIPTION:

ATGCCCCGTG CCCCCCACTC CATGCCCTTG CTGCTGCTGT TGCTGCTGTC ACTCCCCCAA 60
GGCCAGACTG CCTTTCCCA GGACCCATC CCTCTGTGA CCTCTGACCT ACAAGGTACC 120
TCTCCGTCAT CCTGGTTCG GGGCCTGGAG GACGATGCTG TGGCTGCGGA ACTTGGGCTG 180
GACTTTCAGA GATTCTGAC CTTGAACCGG ACCTTGCTTG TGGCTGCCCG GGATCACGTT 240

TTCTCCTTCG ATCTTCAAGC CCAAGAAGAA GGGGAGGGGC TGGTGOCCAA CAAGTTTCTG 300
 5 ACATGGCGGA GCCAAGACAT GGAGAATTGT GCTGTCCGGG GAAAGCTGAC GGACGAATGC 360
 TACAACCTACA TCCGTGTTCT TGTTCCTGG GACTCGCAGA CACTCCTTGC CTGTGGAACA 420
 AATTCCTTCA GCCCTGTGTG TCGCAGCTAT GGGATAACAT CTCTGCAACA GGAGGGTGAG 480
 10 GAGCTGAGTG GGCAAGCTCG ATGCCCTTT GATGCCAACC AGTCCACTGT GGCCATCTCT 540
 GCAGAGGGTA GTTTGTAATC AGCCACAGCA GCAGATTTC AGGCCAGTGA TGCTGTGGTT 600
 TACAGAAGCC TTGGACCTCA GCCCCCACTC CGTTCTGCAA AGTATGACTC CAAGTGGCTT 660
 15 CGAGAGCCAC ACTTTGTCTA TGCTTTGGAG CATGGAGACC ATGTCTACTT CTTTCTCCG 720
 GAGAAGTCTC TGTGGAGGAC GCGCGGCTG GGGAGGGTGC AGTTTCCCG GGTGGCCCGG 780
 20 CTGTGTAAAC GTGACATGGG TGGCTACCA CCGGCCTTG ATGCCACTG GACATCCTTC 840
 CTTAAGCTGA GGCTCAACTG CTCCGTCCCT GGGGACTCTA CCTTCTACTT TGATGTCTTA 900
 25 CAGTCTTAA CTGGGCTGT GAACCTGCAT GGGCGCTCTG CCCTCTTTGG GGTCTTCACT 960
 ACTCAGACCA ATAGCATTCC TGGGTCTGCA GTCTGCGCCT TCTACCTAGA TGACATTGAA 1020
 CGTGGCTTTG AGGGCAAGTT CAAGGAGCAG AGGAGTCTGG ATGGGCGCTG GACTCCTGTG 1080
 30 TCTGAGGACA AAGTCCCTC ACCCAGGCCA GGTCTCTGTG CAGGTGTGGG TGCAGCTGCC 1140
 TTATTCTCCT CCTCTCAAGA CCTGCCTGAC GATGTCCTGC TCTTCATCA GGCACACCA 1200
 35 CTGCTGGATC CCGCTGTGCC ACCTGCCACC CATCAACCTC TCCTACTCT GACTAGCAGG 1260
 GCTCTACTGA CCCAGGTAGC TGTGGATGGT ATGGCTGGCC CCCACAGAAA TACTACAGTC 1320
 CTGTTTCTTG GCTCCAATGA TGGGACAGTG CTGAAGTGC TACCTCCAGG GGGACAGTCT 1380
 40 CTGGGACCCG AGCCTATCAT ATTGAAGAG ATTGATGCCT ACAGCCATGC CCGGTGCAGT 1440
 GGAAGCGGT CACCCGAGC TGCTCGACGG ATCATAGGGC TGGAGCTGGA CACTGAGGGT 1500
 45 CACAGGCTTT TTGTGGCCTT TCCTGGATGC ATCGTCTACC TCTCTCTCAG CCGCTGTGCC 1560
 CGGCATGGAG CATGTCAGAG GAGCTGCTG GCTTCTCTGG ACCCATACTG TGGATGGCAT 1620
 50 CCGTCCGAG GCTGTGTGAA TATCAGGGA CCTGGAGGGA CTGATGTGGA TCTGACTGGG 1680
 AACCAGGAAT CCATGGAGCA TGGTACTGC CAAGATGGAG CGACTGGGAG TCAGTCTGCC 1740

55

5
10
15
20
25
30
35
40
45
50
55

CCTGGAGATT CTGCTATGG CGTGCGCAGG GACCTTTCCC CAGCCTCAGC CTCCCGATCC 1800
ATCCCCATCC CACTCCTCCT GGCCTGTGTG GCGGCGGCGT TCGCTTTGGG CGCCTCAGTC 1860
TCCGGCCTCT TGGTGTCTG TGCTTGTCTG CGCGCGAACC GCCGTGGAG CAAGGACATC 1920
GAGACCCCGG GGCTGCCCG CCCCCTCTCC CTTCGCAGTC TGGCGAGGCT GCACGGTGGC 1980
GGTCTGAGC CCCCCTCTCC GCCAAGGAT GGTGATGAG CGCAAACGCC CCAGCTCTAC 2040
ACTACCTTCC TGCCTCCGCC CGAGGGCGGA TCCCACCGG AGCTGGCCTG CCTGCCACC 2100
CCGGAGACCA CGCCCGAGCT GCCGTGAAG CACCTCCGTG CCTCCGGGG TCCCTGGGAG 2160
TGGAACCAGA ACGGGAACAA CGCTTCGGAG GGCCAGGCC GCCACGGGG CTGCAGCGC 2220
GCGGGCGGGC CCGCCCGCG CGTGCTGGT AGGCCACGC CCCCTGGCTG CCCCAGGAG 2280
GAGGTGGAGG TGACCACGCT GGAGAACTG CTGCGTACC TGCACGGCCC GCAGCCGCC 2340
AGGAAGGCA GGAACCTCT CGCCTCCGCC CCGTTCACCT CCGGCGGCC TGCTCGAG 2400
CCCGCGCGC CTTGTTCGT GGAATCCAGC CCGATGCTC GTGATTGCTG GCCGCCGCTG 2460
AGGCTCGAG TACCGCCGA CGCAAGCGC GCGGCCCGA GCGGCGGCC TGCTCTCTG 2520
GCCCCGGCTC CACGCTGGG CGTCAGCGC AGCCGAAGAT TGCCCTCCC CACGCACCG 2580
GCGCCCCCGG GCTGTCTAC CCGAGTCCC TCGGAGGCC CGTCCAGTA CTCGGGGGG 2640
CCCGGAGGC ACCTCTGTA CCTGGGCGG CCGACGGCC ACCGCGCGC CTCCTGAAG 2700
AGGGTGGAG TGAAGTCTC ACTGTCGCC AAACCGCCC TCGCCACAC GCCGAGCCC 2760
GCCCCGCAG GCAGCCATT TAACCTC 2787

SEQ ID NO: 3

SEQUENCE LENGTH: 929 amino acids

SEQUENCE TYPE: amino acid

TOPOLOGY: linear

MOLECULE TYPE: peptide

ORIGINAL SOURCE:

ORGANISM: rat (*Rattus norvegicus*)

TISSUE TYPE: brain

FEATURE:

FEATURE KEY: peptide

LOCATION: 1..929

IDENTIFICATION METHOD: P

SEQUENCE DESCRIPTION:

Met Pro Arg Ala Pro His Ser Met Pro Leu Leu Leu Leu Leu Leu Leu

5 10 15

Ser Leu Pro Gln Ala Gln Thr Ala Phe Pro Gln Asp Pro Ile Pro Leu

20 25 30

Leu Thr Ser Asp Leu Gln Gly Thr Ser Pro Ser Ser Trp Phe Arg Gly

35 40 45

Leu Glu Asp Asp Ala Val Ala Ala Glu Leu Gly Leu Asp Phe Gln Arg

50 55 60

Phe Leu Thr Leu Asn Arg Thr Leu Leu Val Ala Ala Arg Asp His Val

65 70 75 80

Phe Ser Phe Asp Leu Gln Ala Gln Glu Glu Gly Glu Gly Leu Val Pro

85 90 95

Asn Lys Phe Leu Thr Trp Arg Ser Gln Asp Met Glu Asn Cys Ala Val

100 105 110

Arg Gly Lys Leu Thr Asp Glu Cys Tyr Asn Tyr Ile Arg Val Leu Val

115 120 125

Pro Trp Asp Ser Gln Thr Leu Leu Ala Cys Gly Thr Asn Ser Phe Ser

130 135 140

Pro Val Cys Arg Ser Tyr Gly Ile Thr Ser Leu Gln Gln Glu Gly Glu
 145 150 155 160
 5 Glu Leu Ser Gly Gln Ala Arg Cys Pro Phe Asp Ala Thr Gln Ser Thr
 165 170 175
 10 Val Ala Ile Ser Ala Glu Gly Ser Leu Tyr Ser Ala Thr Ala Ala Asp
 180 185 190
 15 Phe Gln Ala Ser Asp Ala Val Val Tyr Arg Ser Leu Gly Pro Gln Pro
 195 200 205
 20 Pro Leu Arg Ser Ala Lys Tyr Asp Ser Lys Trp Leu Arg Glu Pro His
 210 215 220
 Phe Val Tyr Ala Leu Glu His Gly Asp His Val Tyr Phe Phe Leu Pro
 25 225 230 235 240
 Glu Lys Ser Leu Trp Arg Thr Pro Gly Leu Gly Arg Val Gln Phe Ser
 245 250 255
 30 Arg Val Ala Arg Val Cys Lys Arg Asp Met Gly Gly Ser Pro Arg Ala
 260 265 270
 35 Leu Asp Arg His Trp Thr Ser Phe Leu Lys Leu Arg Leu Asn Cys Ser
 275 280 285
 40 Val Pro Gly Asp Ser Thr Phe Tyr Phe Asp Val Leu Gln Ser Leu Thr
 290 295 300
 Gly Pro Val Asn Leu His Gly Arg Ser Ala Leu Phe Gly Val Phe Thr
 45 305 310 315 320
 Thr Gln Thr Asn Ser Ile Pro Gly Ser Ala Val Cys Ala Phe Tyr Leu
 325 330 335
 50 Asp Asp Ile Glu Arg Gly Phe Glu Gly Lys Phe Lys Glu Gln Arg Ser
 55

340 345 350
 Leu Asp Gly Ala Trp Thr Pro Val Ser Glu Asp Lys Val Pro Ser Pro
 5 355 360 365
 Arg Pro Gly Ser Cys Ala Gly Val Gly Ala Ala Ala Leu Phe Ser Ser
 10 370 375 380
 Ser Gln Asp Leu Pro Asp Asp Val Leu Leu Phe Ile Lys Ala His Pro
 15 385 390 395 400
 Leu Leu Asp Pro Ala Val Pro Pro Ala Thr His Gln Pro Leu Leu Thr
 20 405 410 415
 Leu Thr Ser Arg Ala Leu Leu Thr Gln Val Ala Val Asp Gly Met Ala
 25 420 425 430
 Gly Pro His Arg Asn Thr Thr Val Leu Phe Leu Gly Ser Asn Asp Gly
 30 435 440 445
 Thr Val Leu Lys Val Leu Pro Pro Gly Gly Gln Ser Leu Gly Pro Glu
 35 450 455 460
 Pro Ile Ile Leu Glu Glu Ile Asp Ala Tyr Ser His Ala Arg Cys Ser
 40 465 470 475 480
 Gly Lys Arg Ser Pro Arg Ala Ala Arg Arg Ile Ile Gly Leu Glu Leu
 45 485 490 495
 Asp Thr Glu Gly His Arg Leu Phe Val Ala Phe Pro Gly Cys Ile Val
 50 500 505 510
 Tyr Leu Ser Leu Ser Arg Cys Ala Arg His Gly Ala Cys Gln Arg Ser
 55 515 520 525
 Cys Leu Ala Ser Leu Asp Pro Tyr Cys Gly Trp His Arg Phe Arg Gly
 530 535 540

Cys Val Asn Ile Arg Gly Pro Gly Gly Thr Asp Val Asp Leu Thr Gly
 545 550 555 560
 5 Asn Gln Glu Ser Met Glu His Gly Asp Cys Gln Asp Gly Ala Thr Gly
 565 570 575
 10 Ser Gln Ser Gly Pro Gly Asp Ser Ala Tyr Gly Val Arg Arg Asp Leu
 580 585 590
 15 Ser Pro Ala Ser Ala Ser Arg Ser Ile Pro Ile Pro Leu Leu Leu Ala
 595 600 605
 20 Cys Val Ala Ala Ala Phe Ala Leu Gly Ala Ser Val Ser Gly Leu Leu
 610 615 620
 Val Ser Cys Ala Cys Arg Arg Ala Asn Arg Arg Arg Ser Lys Asp Ile
 625 630 635 640
 25 Glu Thr Pro Gly Leu Pro Arg Pro Leu Ser Leu Arg Ser Leu Ala Arg
 645 650 655
 30 Leu His Gly Gly Gly Pro Glu Pro Pro Pro Pro Pro Lys Asp Gly Asp
 660 665 670
 35 Ala Ala Gln Thr Pro Gln Leu Tyr Thr Thr Phe Leu Pro Pro Pro Glu
 675 680 685
 40 Gly Gly Ser Pro Pro Glu Leu Ala Cys Leu Pro Thr Pro Glu Thr Thr
 690 695 700
 Pro Glu Leu Pro Val Lys His Leu Arg Ala Ser Gly Gly Pro Trp Glu
 705 710 715 720
 45 Trp Asn Gln Asn Gly Asn Asn Ala Ser Glu Gly Pro Gly Arg Pro Arg
 725 730 735
 50 Gly Cys Ser Ala Ala Gly Gly Pro Ala Pro Arg Val Leu Val Arg Pro
 55

	740	745	750
	Pro Pro Pro Gly Cys Pro Gly Gln Glu Val Glu Val Thr Thr Leu Glu		
5	755	760	765
	Glu Leu Leu Arg Tyr Leu His Gly Pro Gln Pro Pro Arg Lys Gly Ser		
10	770	775	780
	Glu Pro Leu Ala Ser Ala Pro Phe Thr Ser Arg Pro Pro Ala Ser Glu		
15	785	790	795
	Pro Gly Ala Ala Leu Phe Val Asp Ser Ser Pro Met Pro Arg Asp Cys		
20	805	810	815
	Val Pro Pro Leu Arg Leu Asp Val Pro Pro Asp Gly Lys Arg Ala Ala		
25	820	825	830
	Pro Ser Gly Arg Pro Ala Leu Ser Ala Pro Ala Pro Arg Leu Gly Val		
30	835	840	845
	Ser Gly Ser Arg Arg Leu Pro Phe Pro Thr His Arg Ala Pro Pro Gly		
35	850	855	860
	Leu Leu Thr Arg Val Pro Ser Gly Gly Pro Ser Arg Tyr Ser Gly Gly		
40	865	870	875
	Pro Gly Arg His Leu Leu Tyr Leu Gly Arg Pro Asp Gly His Arg Gly		
45	885	890	895
	Arg Ser Leu Lys Arg Val Asp Val Lys Ser Pro Leu Ser Pro Lys Pro		
50	900	905	910
	Pro Leu Ala Thr Pro Pro Gln Pro Ala Pro His Gly Ser His Phe Asn		
55	915	920	925
	Phe		

SEQ ID NO: 4

SEQUENCE LENGTH: 3432 base pairs

SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

TOPOLOGY: linear

MOLECULE TYPE: cDNA to mRNA

HYPOTHETICAL: No

ANTI-SENSE: No

ORIGINAL SOURCE:

ORGANISM: human (Homo sapiens)

TISSUE TYPE: child brain

FEATURE:

FEATURE KEY: 5' UTR

LOCATION: 1..187

IDENTIFICATION METHOD: E

FEATURE KEY: CDS

LOCATION: 188..2977

IDENTIFICATION METHOD: E

FEATURE KEY: 3' UTR

LOCATION: 2978..3407

IDENTIFICATION METHOD: E

FEATURE KEY: polyA signal

LOCATION: 3408..3432

IDENTIFICATION METHOD: E

SEQUENCE DESCRIPTION:

5
10
15
20
25
30
35
40
45
50
55

AAAACCACGG ATTGCGAACT CAGCGCAGCG CGTGGCCGCT GGC CGCCCGC GGCGATCTCG 60
ATCCCGCTGA CCCGAATCCT GGAGTCAGAG GTTTCCTATC CCCCTCAAGC CCCACAGGA 120
GTCACCAACC CAGGGCCGGC TTATGGGTGA GGGGGCACCC CCTGGGGCCT GAGCTGCCCC 180
ACACAGGATG CCCCGTGCCC CCCACTTCAT GCCCTTGCTG CTA CTGCTGC TGCTGCTCTC 240
ACTTCCCAT ACTCAGGCGG CCTTCCCCA GGACCCCTC CCTCTGTGA TCTCTGACCT 300
TCAAGGTACT TCCCCATTAT CTG GTTTCG GGGCCTGGAG GATGATGCTG TGGCTGCAGA 360
ACTTGGGCTG GACTTTCAGA GATTCTGAC CTTGAACCGG ACCTTGCTAG TGGCTGCCCG 420
GGATCACGTT TTCTCCTCG ATCTTCAAGC CGAAGAAGAA GGGGAGGGG TGGTGCCAA 480
CAAGTATCTA ACATGGAGAA GCCAAGATGT GGAGAACTGT GCTGTACGGG GAAAGCTGAC 540
GGATGAGTGC TACAACTATA TTCGTGTCT TGTCCCTGG GACTCCAGA CGCTCCTTG 600
CTGTGGAACG AACTCATTCA GCCCTGTGT CCGCAGCTAT GGGATAACTT CGCTGCAGCA 660
GGAGGGTGAG GAACTGAGTG GGCAGGCTCG ATGCCCTTT GATGCCACCC AGTCCAACGT 720
GGCCATCTTT CGAGAGGGCA GCCTGTACTC AGCCACAGCT GCGGATTTC AGGCCAGTGA 780
TGCTGTAGTT TACAGAAGCC TTGGGCCCCA GCCCCACTC CGCTCCGCCA AGTATGACTC 840
CAAGTGGCTC CGAGAGCCAC ACTTTGTCCA GGCCTTGGAG CATGGAGACC ATGTCTACTT 900
CTTCTTCCG GAGGTCTCTG TGGAGGATGC TCGGCTGGG AAGGTGCAGT TCTCCCGCGT 960
AGCCCGAGTA TGTAACGTG ACATGGGCGG CTCGCCCTCG GCCTTGACC GCCACTGGAC 1020
ATCCTTCCTG AAGCTTCGGC TCAACTGCTC TGTCCCTGG GACTCTACTT TCTATTTGA 1080
TGTTTTACAG GCCTTGACTG GGCTGTGAA CCTGCATGGC CGCTCTGCTC TCTTTGGGT 1140
CTTACCACC CAGACCAATA GCATCCCTGG CTCTGCCGTC TGCGCCTTCT ACCTGGATGA 1200
GATTGAGCGT GGGTTTGAGG GCAAGTTCAA GGAGCAGAG AGTCTGGATG GGGCCTGGAC 1260
TCCTGTGTCT GAGGACAGAG TTCCCTCACC CAGGCCAGGA TCCTGTGCAG GAGTAGGGG 1320

AGCTGCCTTG TTCTCTCTT CCCGAGACCT CCCTGATGAT GTCCTGACCT TCATCAAGGC 1380
TCACCCGCTG CTGGACCCCG CTGTACCACC TGTACCCAT CAGCCTCTAC TCACTCTCAC 1440
5 TAGCAGGGCC CTACTGACCC AAGTAGCTGT GGATGGCATG GCTGGTCCC ACAGTAACAT 1500
CACAGTCATG TTCTTGGCT CCAATGATGG GACAGTCTG AAGGTGCTGA CCCAGGTGG 1560
10 GCGATCCGGG GGACCTGAGC CCATCCTCT GGAAGAGATT GATGCCTACA GCCCTGCCCG 1620
GTGCAGTGGG AAGCGGACAG CCCAAACAGC ACGACGGATC ATAGGGCTGG AGCTGGACAC 1680
TGAGGGTCAC AGGCTTTTTG TGGCTTTTTC TGGCTGTATT GTCTACCTCC CTCTCAGCCG 1740
15 GTGTGCCCGG CATGGGCCCT GTCAGAGGAG CTGTTTGGCT TCTCAGGACC CATACTGTGG 1800
ATGGCATAGC TCCAGGGGCT GTGTGGATAT CAGGGGATCT GGTGGGACTG ATGTGGATCA 1860
20 GGCTGGGAAC CAGGAATCCA TGGAGCATGG TGAAGGCTCA GATGGAGCTA CTGGGAGTCA 1920
GTCTGGCCCT GGGGATCTG CTATGGCGT GCGCCGGGAC CTGCCCCAG CCTCGGCTC 1980
CCGCTCCGTC CCATCCAC TCCTCTGGC CAGTGTGGC GCAGCTTTG CCTGGGCGC 2040
25 CTGAGTCTCT GGCTCTGG TCTCTGTGC TTGTGCGCG GCCACCGAC GTCGGGGCAA 2100
GGACATCGAG ACTCCGGGG TCCCGGCCC TCTCTCCCTC CGCAGTTTG CCCGGCTCCA 2160
30 CCGTGGGGG CCAGAGCCCC CGCCGCCCTC CAAGGACGGG GACGCGTGC AGACGCCGCA 2220
GCTCTACAC ACCTTCTGC CGCTCCGGA GGGCGTCCC CGCCGGAGC TGGCTGCT 2280
35 GCCACCCCC GAGTCCACG CGGAGTGCC GGTCAAGCAC CTCGCGCGC CCGGGACCC 2340
CTGGGAGTGG AACCAGAACA GGAACAACG CAAGGAGGT CCGGGCGCT CAGGGGCGG 2400
GCACGCGCG GCGGGGCGG CGCCCCCGT GCTGGTGGG CCACCGCGC CCGGCTGTCC 2460
40 CCGGCAGGCC GTGGAAGTCA CCACCTGGA GGAAGTCTG CGTACCTGC ACGGCCCGCA 2520
GCGGCCAGA AAGGGGCGG AGCCCCCGC CCTTTAACC TCGGGGCGC TCCGCGGA 2580
45 GCGGCCCCG GCGCTCTTG GCGGCCAG CCCCAGGCC CAGAGTGG CCTCGCGCT 2640
GAGGCTGGAC GTGCCCCCG AGGGCAGTG CGCTCTGCC CCGGCCGGC CCGGCTCTC 2700
50 CGCCCCGCT CCGGGCTGG GGTGCGCG AGGCCGAGG TTGCTTTCT CCGGCCACG 2760
GGCCCCCCT GCCCTGCTA CTCAGTCCC CTCGGAGGT CCTCCAGT ACTCGGGG 2820

5 TCCCGGAAG CACCTCTGT ACCTGGGCG GCGGAGGGC TACCGGGCC GCGCCCTGAA 2880
AAGGGTGGAC GTCGAGAAG CCCAGTTGTC CCTGAAGCT CCCCTGTCG GGCCTCCTC 2940
CGCCAGGGC GTCCGAACG GCGGCGTTT CAACTTTAA AGGAGCGGT CCACGGCTC 3000
CAGCGTGGG AGCGCCGAG TCCTCTCGT CACGAGCTG ACGCTCTCA GGACGTTCA 3060
10 CCGCCCTC GCGCGCAC TCCAGCTTC CCGACTCGA GAGTCTCCG AGGCCCTTT 3120
TCGCTCGG TTTATTTATT GACTGTCTT CCCCCTGTC TCGACAGAAG AGTGGGAGT 3180
GAGAAGCCG TCTCTCAGT GAGCCAGCAT TTCAGGGGA GCTGGCGGAC TCCACTCCC 3240
CGCTCCCTC CAGCCAAGT GCCTTAATC GCGCTCGG GCTCCCCAG AGACTGTGCC 3300
CGGGCGGGC CGCGCGGCT GTGTCCAGAG TCCTCGGCC TCCTGGTCT GGGACGTGCC 3360
20 TCTCTACTG TGTAGGAGC TCGCTTCCC AATACAGCG TGTCTGCAA AAAAAAAAAA 3420
AAAAAAAA AA 3432

25 SEQ ID NO: 5

SEQUENCE LENGTH: 2790 base pairs

30 SEQUENCE TYPE: nucleic acid

STRANDEDNESS: double

35 TOPOLOGY: linear

MOLECULE TYPE: cDNA to mRNA

40 HYPOTHETICAL: No

ANTI-SENSE: No

ORIGINAL SOURCE:

45 ORGANISM: human (Homo sapiens)

TISSUE TYPE: child brain

50 FEATURE:

FEATURE KEY: CDS

55

LOCATION: 1..2790

IDENTIFICATION METHOD: E

SEQUENCE DESCRIPTION:

5
10
15
20
25
30
35
40
45
50
55

ATGCCCCGTG CCCCCCACTT CATGCCCTTG CTGCTACTGC TGCTGCTGCT CTCACTTCCC 60
CATACTCAGG CCGCCTTTCC CCAGGACCCC CTCCTCTGT TGATCTCTGA CCTTCAAGGT 120
ACTTCCCAT TATCCTGGTT TCGGGGCTG GAGGATGATG CTGTGGCTGC AGAACTTGGG 180
CTGGACTTTC AGAGATTCT GACCTGAAC CGGACCTGC TAGTGGCTGC CCGGGATCAC 240
GTTTTCTCT TCGATCTTCA AGCCGAAGAA GAAGGGGAGG GGCTGGTGCC CAACAAGTAT 300
CTAACATGGA GAAGCCAAGA TGTGGAGAAC TGTGCTGTAC GGGGAAAGCT GACGGATGAG 360
TGCTACAACT ATATTCTGT TCTTGTCCC TGGGACTCCC AGACGCTCCT TGCTGTGGA 420
ACGAACTCAT TCAGCCCTGT GTGCGCAGC TATGGGATAA CTTGCTGCA GCAGGAGGT 480
GAGGAACTGA GTGGGCAGGC TCGATGCCCC TTTGATGCCA CCCAGTCAA CGTGGCCATC 540
TTTGACAGAG GCAGCCTGTA CTCAGCCACA GCTCCGATT TCCAGGCCAG TGATGCTGTA 600
GTTTACAGAA GCCTTGGGCC CCAGCCCCCA CTCGCTCCG CCAAGTATGA CTCCAAGTGG 660
CTCCGAGAGC CACACTTGT CCAGGCCTTG GAGCATGGAG ACCATGTCTA CTTCTTCTT 720
CGCGAGGTCT CTGTGGAGGA TGCTCGGCTG GGGAAAGTGC AGTTCTCCCG CGTAGCCCGA 780
GTATGTAAAC GTGACATGGG CGGCTCGCCT CGGGCCTTG ACCGCCACTG GACATCCTT 840
CTGAAGCTTC GGCTCAACTG CTCTGTCCCT GGGGACTCTA CTTTCTATTT TGATGTTTTA 900
CAGGCCTTGA CTGGGCTGT GAACCTGCAT GGCCGCTCTG CTCTCTTTGG GGTCTTCACC 960
ACCCAGACCA ATAGCATCCC TGGCTCTGCC GTCTGCGCCT TCTACCTGGA TGAGATTGAG 1020
CGTGGGTTTG AGGGCAAGTT CAAGGAGCAG AGGAGTCTGG ATGGGGCCTG GACTCCTGTG 1080
TCTGAGGACA GAGTTCCCTC ACCCAGGCCA GGATCCTGTG CAGGAGTAGG GGGAGCTGCC 1140
TTGTTCTCCT CTTCCCGAGA CCTCCCTGAT GATGTCTGA CTTTCATCA GGCTCACCCG 1200
CTGCTGGACC CCGCTGTACC ACCTGTACCC CATCAGCCTC TACTACTCT CACTAGCAGG 1260
GCCCTACTGA CCCAAGTAGC TGTGGATGGC ATGGCTGGTC CCCACAGTAA CATCACAGTC 1320

5
10
15
20
25
30
35
40
45
50
55

ATGTTCCCTG GCTCCAATGA TGGGACAGTG CTGAAGGTGC TGACCCAGG TGGGCGATCC 1380
GGGGGACCTG AGCCCATCCT CCTGGAAGAG ATTGATGCCT ACAGCCCTGC CCGTGCACT 1440
GGGAAGCGGA CAGCCCAAAC AGCACGACGG ATCATAGGCC TGGAGCTGGA CACTGAGGGT 1500
CACAGGCTTT TTGTGGCTTT TTCTGGCTGT ATTGTCTACC TCCCTCTCAG CCGGTGTGCC 1560
CGGCATGGGG CCTGTCAGAG GAGCTGTTTG GCTTCTCAGG ACCATACTG TGGATGGCAT 1620
AGCTCCAGGG GCTGTGTGGA TATCAGGGGA TCTGGTGGGA CTGATGTGGA TCAGGCTGGG 1680
AACCAGGAAT CCATGGAGCA TGGTACTGC CAAGATGGAG CTAAGGGAG TCAGTCTGGC 1740
CCTGGGGATT CTGCTTATGG CGTGCCCGG GACCTGCCCC CAGCCTCGGC CTCCCGCTCC 1800
GTCCCCATCC CACTCCTCCT GGCCAGTGTG GCCGAGCTT TTGCCCTGGG CGCCTCAGTC 1860
TCTGGCCTCC TGGTCTCCTG TGCTTGTGCG CGGCCCCACC GACGTGGGG CAAGGACATC 1920
GAGACTCCCG GGCTCCCGCG CCTCTCTCC CTCCGAGTT TGGCCCGCT CCACGGTGGG 1980
GGCCAGAGC CCGCGCCGC CTCCAAGGAC GGGGACCGG TGCAGACGCC GCAGCTCTAC 2040
ACCACCTTCC TGCCGCTCC GGAGGGCGTG CCCCCCGCG AGCTGGCTG CCTGCCACC 2100
CCCGAGTCCA CGCCGAGCT GCCGTCAAG CACCTCCGG CGCCGGGGA CCCCTGGGAG 2160
TGGAACCAGA ACAGGAACAA CGCCAAGGAG GGTCCGGCC GCTACGGGG CGGGCAGCG 2220
CGGGGCGGG CGCGCCCCG CGTGTGGTG AGGCCACCG CGCCGGCTG TCOCGGGAG 2280
GCCGTGGAAG TCACCACCCT GGAGGAACTG CTGCGTACC TGCACGGCC GCAGCCGCC 2340
AGAAAGGGG CCGAGCCCC CGCCCTTTA ACCTCGCGG CGCTCCGCC GGAGCCGCC 2400
CCCGCCTCT TGGGCGGCC CAGCCCAGG CCCACGAGT GCGCTCGCC GCTGAGGCTG 2460
GACGTGCCCC CCGAGGGCAG GTGCGCTCT GCCCCCGCC GGGCGGCT CTCCGCCCC 2520
GCTCCCCGG TGGGCGTCG CGGAGGCCG AGGTTGCTT TCTCGGCCA CCGGGCCCC 2580
CCTGCCCTG TCACTCGAGT CCCCTCGGA GGTCCCTCA GGTACTCGG GGTCCCGG 2640
AAGCACTCC TGTACTGGG CCGGCCGAG GGCTACGGG GCGCGCCCT GAAAAGGGT 2700
GACGTGAGA AGCCCGAGT GTCCCTGAAG CCTCCCTCG TCGGGCCCTC CTCCGCCAG 2760
GCCGTCCGA ACGGCGCGG TTTCACTTT 2790

SEQ ID NO: 6

SEQUENCE LENGTH: 930 amino acids

SEQUENCE TYPE: amino acid

TOPOLOGY: linear

MOLECULE TYPE: peptide

ORIGINAL SOURCE:

ORGANISM: human (Homo sapiens)

TISSUE TYPE: child brain

FEATURE:

FEATURE KEY: peptide

LOCATION: 1..930

IDENTIFICATION METHOD: P

SEQUENCE DESCRIPTION:

Met Pro Arg Ala Pro His Phe Met Pro Leu Leu Leu Leu Leu Leu

5 10 15

Leu Ser Leu Pro His Thr Gln Ala Ala Phe Pro Gln Asp Pro Leu Pro

20 25 30

Leu Leu Ile Ser Asp Leu Gln Gly Thr Ser Pro Leu Ser Trp Phe Arg

35 40 45

Gly Leu Glu Asp Asp Ala Val Ala Ala Glu Leu Gly Leu Asp Phe Gln

50 55 60

Arg Phe Leu Thr Leu Asn Arg Thr Leu Leu Val Ala Ala Arg Asp His

65 70 75 80

Val Phe Ser Phe Asp Leu Gln Ala Glu Glu Glu Gly Glu Gly Leu Val

	85	90	95
	Pro Asn Lys Tyr Leu Thr Trp Arg Ser Gln Asp Val Glu Asn Cys Ala		
5	100	105	110
	Val Arg Gly Lys Leu Thr Asp Glu Cys Tyr Asn Tyr Ile Arg Val Leu		
10	115	120	125
	Val Pro Trp Asp Ser Gln Thr Leu Leu Ala Cys Gly Thr Asn Ser Phe		
15	130	135	140
	Ser Pro Val Cys Arg Ser Tyr Gly Ile Thr Ser Leu Gln Gln Glu Gly		
	145	150	155
20	Glu Glu Leu Ser Gly Gln Ala Arg Cys Pro Phe Asp Ala Thr Gln Ser		
	165	170	175
25	Asn Val Ala Ile Phe Ala Glu Gly Ser Leu Tyr Ser Ala Thr Ala Ala		
	180	185	190
	Asp Phe Gln Ala Ser Asp Ala Val Val Tyr Arg Ser Leu Gly Pro Gln		
30	195	200	205
	Pro Pro Leu Arg Ser Ala Lys Tyr Asp Ser Lys Trp Leu Arg Glu Pro		
35	210	215	220
	His Phe Val Gln Ala Leu Glu His Gly Asp His Val Tyr Phe Phe Phe		
40	225	230	235
	Arg Glu Val Ser Val Glu Asp Ala Arg Leu Gly Lys Val Gln Phe Ser		
	245	250	255
45	Arg Val Ala Arg Val Cys Lys Arg Asp Met Gly Gly Ser Pro Arg Ala		
	260	265	270
50	Leu Asp Arg His Trp Thr Ser Phe Leu Lys Leu Arg Leu Asn Cys Ser		
	275	280	285

55

Val Pro Gly Asp Ser Thr Phe Tyr Phe Asp Val Leu Gln Ala Leu Thr
 290 295 300
 5 Gly Pro Val Asn Leu His Gly Arg Ser Ala Leu Phe Gly Val Phe Thr
 305 310 315 320
 10 Thr Gln Thr Asn Ser Ile Pro Gly Ser Ala Val Cys Ala Phe Tyr Leu
 325 330 335
 15 Asp Glu Ile Glu Arg Gly Phe Glu Gly Lys Phe Lys Glu Gln Arg Ser
 340 345 350
 20 Leu Asp Gly Ala Trp Thr Pro Val Ser Glu Asp Arg Val Pro Ser Pro
 355 360 365
 Arg Pro Gly Ser Cys Ala Gly Val Gly Gly Ala Ala Leu Phe Ser Ser
 25 370 375 380
 Ser Arg Asp Leu Pro Asp Asp Val Leu Thr Phe Ile Lys Ala His Pro
 385 390 395 400
 30 Leu Leu Asp Pro Ala Val Pro Pro Val Thr His Gln Pro Leu Leu Thr
 405 410 415
 35 Leu Thr Ser Arg Ala Leu Leu Thr Gln Val Ala Val Asp Gly Met Ala
 420 425 430
 40 Gly Pro His Ser Asn Ile Thr Val Met Phe Leu Gly Ser Asn Asp Gly
 435 440 445
 45 Thr Val Leu Lys Val Leu Thr Pro Gly Gly Arg Ser Gly Gly Pro Glu
 450 455 460
 Pro Ile Leu Leu Glu Glu Ile Asp Ala Tyr Ser Pro Ala Arg Cys Ser
 465 470 475 480
 50 Gly Lys Arg Thr Ala Gln Thr Ala Arg Arg Ile Ile Gly Leu Glu Leu
 55

EP 0 960 937 A1

	485	490	495
5	Asp Thr Glu Gly His Arg Leu Phe Val Ala Phe Ser Gly Cys Ile Val		
	500	505	510
	Tyr Leu Pro Leu Ser Arg Cys Ala Arg His Gly Ala Cys Gln Arg Ser		
10	515	520	525
	Cys Leu Ala Ser Gln Asp Pro Tyr Cys Gly Trp His Ser Ser Arg Gly		
15	530	535	540
	Cys Val Asp Ile Arg Gly Ser Gly Gly Thr Asp Val Asp Gln Ala Gly		
	545	550	555
20	Asn Gln Glu Ser Met Glu His Gly Asp Cys Gln Asp Gly Ala Thr Gly		
	565	570	575
25	Ser Gln Ser Gly Pro Gly Asp Ser Ala Tyr Gly Val Arg Arg Asp Leu		
	580	585	590
	Pro Pro Ala Ser Ala Ser Arg Ser Val Pro Ile Pro Leu Leu Leu Ala		
30	595	600	605
	Ser Val Ala Ala Ala Phe Ala Leu Gly Ala Ser Val Ser Gly Leu Leu		
35	610	615	620
	Val Ser Cys Ala Cys Arg Arg Ala His Arg Arg Arg Gly Lys Asp Ile		
40	625	630	635
	Glu Thr Pro Gly Leu Pro Arg Pro Leu Ser Leu Arg Ser Leu Ala Arg		
	645	650	655
45	Leu His Gly Gly Gly Pro Glu Pro Pro Pro Pro Ser Lys Asp Gly Asp		
	660	665	670
50	Ala Val Gln Thr Pro Gln Leu Tyr Thr Thr Phe Leu Pro Pro Pro Glu		
	675	680	685

55

Gly Val Pro Pro Pro Glu Leu Ala Cys Leu Pro Thr Pro Glu Ser Thr
 690 695 700
 5 Pro Glu Leu Pro Val Lys His Leu Arg Ala Ala Gly Asp Pro Trp Glu
 705 710 715 720
 10 Trp Asn Gln Asn Arg Asn Asn Ala Lys Glu Gly Pro Gly Arg Ser Arg
 725 730 735
 15 Gly Gly His Ala Ala Gly Gly Pro Ala Pro Arg Val Leu Val Arg Pro
 740 745 750
 20 Pro Pro Pro Gly Cys Pro Gly Gln Ala Val Glu Val Thr Thr Leu Glu
 755 760 765
 25 Glu Leu Leu Arg Tyr Leu His Gly Pro Gln Pro Pro Arg Lys Gly Ala
 770 775 780
 30 Glu Pro Pro Ala Pro Leu Thr Ser Arg Ala Leu Pro Pro Glu Pro Ala
 785 790 795 800
 35 Pro Ala Leu Leu Gly Gly Pro Ser Pro Arg Pro His Glu Cys Ala Ser
 805 810 815
 40 Pro Leu Arg Leu Asp Val Pro Pro Glu Gly Arg Cys Ala Ser Ala Pro
 820 825 830
 45 Ala Arg Pro Ala Leu Ser Ala Pro Ala Pro Arg Leu Gly Val Gly Gly
 835 840 845
 50 Gly Arg Arg Leu Pro Phe Ser Gly His Arg Ala Pro Pro Ala Leu Leu
 850 855 860
 55 Thr Arg Val Pro Ser Gly Gly Pro Ser Arg Tyr Ser Gly Gly Pro Gly
 865 870 875 880
 Lys His Leu Leu Tyr Leu Gly Arg Pro Glu Gly Tyr Arg Gly Arg Ala



20

45

50

55

CTGTTTGGCT TCTCAGGACC CATACTGTGG ATGGCATAGC TCCAGGGGCT GTGTGGATAT 120
 CAGGGGATCT GGTGGGACTG ATGTGGATCA GGCTNGGAAC CAGGAATCCA 170

5

SEQ ID NO: 8

10

SEQUENCE LENGTH: 18 base pairs

SEQUENCE TYPE: nucleic acid

15

STRANDEDNESS: single

TOPOLOGY: linear

MOLECULE TYPE: Other nucleic acid, synthetic DNA

20

HYPOTHETICAL: No

ANTI-SENSE: No

25

FEATURE:

FEATURE KEY: CDS

LOCATION: 1..18

30

IDENTIFICATION METHOD: P

SEQUENCE DESCRIPTION:

35

TGGCTGTATT GTCTACCT

18

40

SEQ ID NO: 9

SEQUENCE LENGTH: 20 base pairs

SEQUENCE TYPE: nucleic acid

45

STRANDEDNESS: single

TOPOLOGY: linear

50

MOLECULE TYPE: Other nucleic acid, synthetic DNA

HYPOTHETICAL: No

55

ANTI-SENSE: Yes

FEATURE:

FEATURE KEY: CDS

LOCATION: 1..20

IDENTIFICATION METHOD: P

SEQUENCE DESCRIPTION:

TGGATTCTG GTTCNAGCC

20

SEQ ID NO: 10

SEQUENCE LENGTH: 18 base pairs

SEQUENCE TYPE: nucleic acid

STRANDEDNESS: single

TOPOLOGY: linear

MOLECULE TYPE: Other nucleic acid, synthetic DNA

HYPOTHETICAL: No

ANTI-SENSE: No

FEATURE:

FEATURE KEY: CDS

LOCATION: 1..18

IDENTIFICATION METHOD: E

SEQUENCE DESCRIPTION:

TGTGTAAACG TGACATGG

18

SEQ ID NO: 11

SEQUENCE LENGTH: 18 base pairs

SEQUENCE TYPE: nucleic acid

STRANDEDNESS: single

TOPOLOGY: linear

MOLECULE TYPE: Other nucleic acid, synthetic DNA

HYPOTHETICAL: No

ANTI-SENSE: Yes

FEATURE:

FEATURE KEY: CDS

LOCATION: 1..18

IDENTIFICATION METHOD: E

SEQUENCE DESCRIPTION:

TGCTAGTCAG AGTGAGGA

18

Claims

1. A gene encoding the following protein (a) or (b):

- (a) Semaphorin Y protein comprising the amino acid sequence shown in SEQ ID NO: 3 or 6,
- (b) a protein which comprises an amino acid sequence wherein one or more amino acids are deleted, substituted and/or added in the amino acid sequence shown in SEQ ID NO: 3 or 6, and which protein inhibits neurite outgrowth.

2. A gene comprising the following DNA (a) or (b):

- (a) Semaphorin Y DNA comprising the base sequence shown in SEQ ID NO: 1, 2, 4, or 5,
- (b) DNA which hybridizes under stringent conditions to DNA comprising the base sequence shown in SEQ ID NO: 1, 2, 4, or 5, and which encodes a protein inhibiting neurite outgrowth.

3. A gene comprising DNA which hybridizes under stringent conditions to DNA comprising the base sequence shown in SEQ ID NO: 7, and which encodes a protein having a semaphorin domain.

4. A protein obtained by expressing the gene of any one of claims 1 to 3.

5. A gene comprising DNA which encodes a protein comprising an amino acid sequence in which one or more amino acids are deleted, substituted and/or added in the protein shown in SEQ ID NO: 3 or 6, wherein said protein promotes neurite outgrowth.

6. A protein obtained by expressing the gene of claim 5.

7. A DNA which is cloned from a human cDNA or genomic library and which hybridizes under stringent conditions to DNA comprising at least part of DNA consisting of the base sequence shown in SEQ ID NO: 1 or 4.
8. An expression plasmid which expresses either the gene of any one of claims 1-3 and 5, or DNA of claim 7.
- 5 9. A transformant transformed with the expression plasmid of claim 8.
- 10 10. A process for producing a recombinant protein, which process comprises culturing the transformant of claim 9, and recovering the recombinant protein expressed.
11. A peptide comprising at least six amino acids of the protein of claim 4 or 6.
12. A peptide of claim 11 which promotes neurite outgrowth.
- 15 13. A peptide of claim 11 characterized in that it contains aspartic acid residue at position 198 of the amino acid sequence shown in SEQ ID NO: 6 or an amino acid residue corresponding to the position of said aspartic acid residue.
- 20 14. An antisense nucleotide, or chemically modified variant thereof, which is directed against a segment comprising at least eight or more bases in the gene of any one of claims 1-3, or in DNA of claim 7.
- 25 15. An antisense nucleotide or chemically modified variant thereof of claim 14, characterized in that it inhibits expression of the protein of claim 4.
- 30 16. An antibody against the protein of claim 4 or 6, or against the peptide of any one of claims 11 to 13.
- 35 17. A pharmaceutical agent comprising, as an active ingredient, the gene of any one of claims 1-3 and 5, DNA of claim 7, the protein of claim 4 or 6, the peptide of any one of claims 11 to 13, the antisense nucleotide or chemically modified variant thereof of claim 14 or 15, or the antibody of claim 16.
- 40 18. A method of screening for Semaphorin Y antagonists, characterized in that it employs the protein of claim 4.
- 45 19. A Semaphorin Y antagonist obtained by the screening method of claim 18.
- 50 20. A Semaphorin Y antagonist of claim 19 which comprises the protein of claim 6, the peptide of any one of claims 11 to 13, or the antibody of claim 16.
- 55 21. A CNS-neuron regeneration promoter, characterized in that it contains at least one of the antisense nucleotides or chemically modified variants thereof of claim 14 or 15, or Semaphorin Y antagonists of claim 19 or 20.
22. A neurite outgrowth inhibitor for PNS-neuron, characterized in that it contains at least one of the proteins of claim 4.
23. A transgenic animal in which either the gene of any one of claims 1-3 and 5, or DNA of claim 7 has been artificially inserted into its chromosome, or has been knocked out.

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP97/03167

A. CLASSIFICATION OF SUBJECT MATTER Int. Cl ⁶ C12N15/12, A61K38/17, A61K31/70, A01K 67/027, C07K14/475 G01N33/577, A61K39/395 // C12P21/08, A61K48/00 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int. Cl ⁶ C12N15/12, A61K38/17, A61K31/70, A01K 67/027, C07H14/475 G01N33/577, A61K39/395, C12P21/08, A61K48/00 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) BIOSIS (DIALOG), WPI (DIALOG), GenBank/EMBL (GENETYX)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	GENOME RESEARCH, 1, 35-42 (1996), G. Lanfranchi et al. "Identification of 4370 Expressed Sequence Tags from a 3'-End-Specific cDNA Library of Human Skeletal Muscle by DNA Sequencing and Filter Hybridization"	7 8-11, 14
A	Cell, Vol. 75, 1389-1399 (1993), Alex L. Kolodkin et al. "The semaphorin Genes Encode a Family of Transmembranand Secreted Growth Cone Guidance Molecules"	1 - 23
A	Neuron, Vol. 14, 941-948 (1995), Andreas W. Puachel et al. "Murine Semaphorin D/Collapsin Is a Member of a Diverse Gene Family and Creates Domains Inhibitory for Axonal Extension"	1 - 23
A	Molecular and Cellular Neuroscience, 9, 26-41 (1997), L. Zhou et al. "Cloning and Expression of a Novel Murine Semaphorin with Structural Similarity to Insect Semaphorin I"	1 - 23
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search December 3, 1997 (03. 12. 97)		Date of mailing of the international search report December 16, 1997 (16. 12. 97)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)